

A new campterochlebiid damselfly dragonfly  
(Odonata: Isophlebioidea) from the Middle Jurassic  
Yanan Formation of Yulin City,  
Shaanxi Province, NW China

Fusheng TIAN, Daran ZHENG, André NEL, Yifei YE,  
Ting MEI & Haichun ZHANG

DIRECTEURS DE LA PUBLICATION / PUBLICATION DIRECTORS :  
Bruno David, Président du Muséum national d'Histoire naturelle  
Étienne Ghys, Secrétaire perpétuel de l'Académie des sciences

RÉDACTEURS EN CHEF / EDITORS-IN-CHIEF: Michel Laurin (CNRS), Philippe Taquet (Académie des sciences)

ASSISTANTE DE RÉDACTION / ASSISTANT EDITOR: Adenise Lopes (Académie des sciences; cr-palevol@academie-sciences.fr)

MISE EN PAGE / PAGE LAYOUT: Audrina Neveu (Muséum national d'Histoire naturelle; audrina.neveu@mnhn.fr)

RÉVISIONS LINGUISTIQUES DES TEXTES ANGLAIS / ENGLISH LANGUAGE REVISIONS: Kevin Padian (University of California at Berkeley)

RÉDACTEURS ASSOCIÉS / ASSOCIATE EDITORS (\*, took charge of the editorial process of the article/a pris en charge le suivi éditorial de l'article):

Micropaléontologie/Micropalaeontology

Lorenzo Consorti (Institute of Marine Sciences, Italian National Research Council, Trieste)

Paléobotanique/Palaeobotany

Cyrille Prestianni (Royal Belgian Institute of Natural Sciences, Brussels)

Métazoaires/Metazoa

**Annalisa Ferretti\*** (Università di Modena e Reggio Emilia, Modena)

Paléochthyologie/Palaeoichthyology

Philippe Janvier (Muséum national d'Histoire naturelle, Académie des sciences, Paris)

Amniotes du Mésozoïque/Mesozoic amniotes

Hans-Dieter Sues (Smithsonian National Museum of Natural History, Washington)

Tortues/Turtles

Walter Joyce (Universität Freiburg, Switzerland)

Lépidosauromorphes/Lepidosauromorphs

Hussam Zaher (Universidade de São Paulo)

Oiseaux/Birds

Eric Buffetaut (CNRS, École Normale Supérieure, Paris)

Paléomammalogie (mammifères de moyenne et grande taille)/Palaeomammalogy (large and mid-sized mammals)

Lorenzo Rook (Università degli Studi di Firenze, Firenze)

Paléomammalogie (petits mammifères sauf Euarchontoglires)/Palaeomammalogy (small mammals except for Euarchontoglires)

Robert Asher (Cambridge University, Cambridge)

Paléomammalogie (Euarchontoglires)/Palaeomammalogy (Euarchontoglires)

K. Christopher Beard (University of Kansas, Lawrence)

Paléoanthropologie/Palaeoanthropology

Aurélien Mounier (CNRS/Muséum national d'Histoire naturelle, Paris)

Archéologie préhistorique (Paléolithique et Mésolithique)/Prehistoric archaeology (Palaeolithic and Mesolithic)

Nicolas Teyssandier (CNRS/Université de Toulouse, Toulouse)

Archéologie préhistorique (Néolithique et âge du bronze)/Prehistoric archaeology (Neolithic and Bronze Age)

Marc Vander Linden (Bournemouth University, Bournemouth)

RÉFÉRÉS / REVIEWERS: <https://sciencepress.mnhn.fr/fr/periodiques/comptes-rendus-palevol/referes-du-journal>

COUVERTURE / COVER:

Made from the Figures of the article.

*Comptes Rendus Palevol* est indexé dans / *Comptes Rendus Palevol is indexed by:*

- Cambridge Scientific Abstracts
- Current Contents® Physical
- Chemical, and Earth Sciences®
- ISI Alerting Services®
- Geoabstracts, Geobase, Georef, Inspec, Pascal
- Science Citation Index®, Science Citation Index Expanded®
- Scopus®.

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Comptes Rendus Palevol* sont référencés par /  
*Articles and nomenclatural novelties published in Comptes Rendus Palevol are registered on:*

- ZooBank® (<http://zoobank.org>)

*Comptes Rendus Palevol* est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris et l'Académie des sciences, Paris  
*Comptes Rendus Palevol is a fast track journal published by the Museum Science Press, Paris and the Académie des sciences, Paris*

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish:*

*Adansonia, Geodiversitas, Zoosystema, Anthropozoologica, European Journal of Taxonomy, Naturae, Cryptogamie* sous-sections *Algologie, Bryologie, Mycologie.*

L'Académie des sciences publie aussi / *The Académie des sciences also publishes:*

*Comptes Rendus Mathématique, Comptes Rendus Physique, Comptes Rendus Mécanique, Comptes Rendus Chimie, Comptes Rendus Géoscience, Comptes Rendus Biologies.*

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél. : 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <https://sciencepress.mnhn.fr>

Académie des sciences, Institut de France, 23 quai de Conti, 75006 Paris.

© This article is licensed under the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>)  
ISSN (imprimé / print): 1631-0683/ ISSN (électronique / electronic): 1777-571X

# A new camptero-phlebiid damsel-dragonfly (Odonata: Isophlebioidea) from the Middle Jurassic Yanan Formation of Yulin City, Shaanxi Province, NW China

**Fusheng TIAN**

Hubei Geological Exploration Institute, China Chemical Geology and Mine Bureau,  
Wuhan 430070 (China)

**Daran ZHENG**

State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology  
and Palaeontology and Center for Excellence in Life and Palaeoenvironment,  
Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008 (China)  
drzheng@nigpas.ac.cn (corresponding author)

**André NEL**

Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle,  
CNRS, Sorbonne Université, EPHE, Université des Antilles,  
case postale 50, 57 rue Cuvier 75231 Paris cedex 05 (France)  
anel@mnhn.fr

**Yifei YE**

Shaanxi Yulin Energy Group Guojiatan Mining Co., Ltd, Yulin 71900 (China)

**Ting MEI**

Hubei Geological Exploration Institute, China Chemical Geology and Mine Bureau,  
Wuhan 430070 (China)

**Haichun ZHANG**

State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology  
and Palaeontology and Center for Excellence in Life and Palaeoenvironment,  
Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008 (China)

Submitted on 8 April 2022 | Accepted on 20 September 2022 | Published on 11 July 2023

---

urn:lsid:zoobank.org:pub:73122322-1744-463F-A812-D4C5BE79A858

---

Tian F., Zheng D., Nel A., Ye Y., Mei T. & Zhang H. 2023. — A new camptero-phlebiid damsel-dragonfly (Odonata: Isophlebioidea) from the Middle Jurassic Yanan Formation of Yulin City, Shaanxi Province, NW China. *Comptes Rendus Palevol* 22 (23): 491–496. <https://doi.org/10.5852/cr-palevol2023v22a23>

## ABSTRACT

The family Camptero-phlebiidae Handlirsch, 1920 is the dominant Jurassic clade of Odonata Fabricius, 1793, especially hosting a high diversification in northern China. The Chinese camptero-phlebiid damsel-dragonflies were mainly recovered from the Middle Jurassic of Inner Mongolia, northern China. In the present study, a new camptero-phlebiid, *Parasinitsia qingyunensis* n. gen., n. sp., is described from the early Middle Jurassic Yanan Formation of the Ordos Basin, NW China.

**KEY WORDS**  
Camptero-  
phlebiidae,  
Yanan Formation,  
Middle Jurassic,  
Ordos Basin,  
China,  
new genus,  
new species.

*Parasinitia* n. gen. resembles the genus *Sinitia* Pritykina, 2006 recorded from the Upper Jurassic of eastern Transbaikalia, but differs from the latter in having two or three rows of cells between RA and RP1 distal of the pterostigma, IR2 and RP3/4 with at least four rows of cells near the wing margin, and CuAa with about 15 rows of cells between it and the posterior wing margin in the broadest area. The new damsel-dragonfly comes from a new early Middle Jurassic insect site from the Ordos Basin, providing new clues to understand the terrestrial ecosystems during this epoch.

**RÉSUMÉ**

*Un nouveau camptero-phlebiid demoiselle-libellule (Odonata: Isophlebioidea) de la formation Yanan du Jurassique moyen de la ville de Yulin, province de Shaanxi, au Nord-Ouest de la Chine.*

La famille Camptero-phlebiidae Handlirsch, 1920 est le clade jurassique dominant des Odonata Fabricius, 1793, avec notamment une forte diversité dans le Nord de la Chine. Les Camptero-phlebiidae de Chine proviennent principalement du Jurassique moyen de la Mongolie intérieure. Dans la présente étude, un nouveau Camptero-phlebiidae, *Parasinitia qingyunensis* n. gen., n. sp., est décrit de la formation de Yanan du Jurassique moyen inférieur du bassin de l'Ordos, dans le nord-ouest de la Chine. *Parasinitia* n. gen. ressemble au genre *Sinitia* Pritykina, 2006 du Jurassique supérieur de l'est de la Transbaikalie, mais diffère de ce dernier en ayant deux ou trois rangées de cellules entre RA et RP1 en aval du pterostigma, IR2 et RP3/4 avec au moins quatre rangées de cellules près du bord de l'aile, et CuAa avec environ 15 rangées de cellules entre elle et la marge postérieure de l'aile dans la zone la plus large. Le nouveau taxon provient d'un nouveau site d'insectes du début du Jurassique moyen du bassin de l'Ordos, fournissant de nouveaux indices pour comprendre les écosystèmes terrestres à cette époque.

**MOTS CLÉS**  
Camptero-  
phlebiidae,  
Formation Yanan,  
Jurassique moyen,  
bassin de l'Ordos,  
Chine,  
genre nouveau,  
espèce nouvelle.

**INTRODUCTION**

The family Camptero-phlebiidae Handlirsch, 1920 is the most diversified damsel-dragonfly clade discovered from the Mesozoic of China, the majority of taxa being recorded from the Middle Jurassic of Inner Mongolia, and few in the Lower Jurassic of Xinjiang and the Middle Jurassic of Shaanxi (Fleck & Nel 2002; Nel *et al.* 2008, 2009; Zheng *et al.* 2016, 2017, 2018, 2019; Huang *et al.* 2018). Herein, we describe a new camptero-phlebiid damsel-dragonfly from the early Middle Jurassic Yanan Formation in Qingyun Village of eastern Yulin City, Shaanxi Province, northwestern China. The Yanan Formation in Huangjiagou Village of Yulin has already yielded a camptero-phlebiid damsel-dragonfly (Zheng *et al.* 2017). The present study describes a new genus and species of Camptero-phlebiidae, and reveals a new Middle Jurassic entomofauna from northern Ordos Basin.

**MATERIAL AND METHODS**

The newly described damsel-dragonfly described was collected from the early Middle Jurassic Yanan Formation in Qingyun Town, Yulin City, Shaanxi Province, NW China (Fig. 1). The Yanan Formation unconformably underlies the Pliocene Baode Formation and conformably overlies the Lower Jurassic Fuxian Formation in Yulin. It is considered to be early Middle Jurassic in age (*c.* 174-172 Ma; Zhang *et al.* 2021). Preserved together with the new damsel-dragonfly, there are abundant fossils including fishes, bivalves, gastropods, plants, and representatives of some other insect orders (Ephemero-

tera Hyatt & Arms, 1891, Phasmatoidea Jacobson & Bianchi, 1902, Coleoptera Linnaeus, 1758, Mecoptera Hyatt & Arms, 1891, Neuroptera Linnaeus, 1758, Orthoptera Latreille, 1793 and Hemiptera Linnaeus, 1758).

The specimen was examined dry using a Nikon SMZ1000 stereomicroscope. Observation was augmented by temporary wetting with laboratory alcohol which improved the contrast between the fossil and the matrix, eliminating the surface irregularity of the latter. Photographs were taken using a Canon 5D digital camera and the line drawings were prepared from photographs using image-editing software (CorelDraw X7 and Adobe Photoshop CS6). The specimen is housed in the Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences (NIGPAS). All taxonomic acts established in the present work have been registered in ZooBank.

The higher classification of fossil and extant Odonatoptera is based on the phylogenetic system of Bechly (1996). The nomenclature of the dragonfly wing venation used in this paper is based on the interpretations of Riek (1976) and Riek & Kukalová-Peck (1984), as modified by Nel *et al.* (1993) and Bechly (1996).

**ABBREVIATIONS**

- AA anal anterior;
- AP anal posterior;
- Arc arculus;
- Ax primary antenodal crossvein;
- C costa;
- Cr nodal crossvein;
- CuAa distal branch of cubitus anterior;
- CuAb proximal branch of cubitus anterior;

CuP	cubitus posterior;
DC	discoidal cell;
IR	intercalary radial vein;
MAa	anterior branch of median anterior;
MAB	posterior branch of median anterior;
MP	median posterior;
N	nodus;
'O'	oblique vein;
Pt	pterostigma;
RA	radius anterior;
RP	radius posterior;
Sn	subnodal crossvein;
ScP	subcosta posterior.

## SYSTEMATIC PALAEOONTOLOGY

Order ODNATA Fabricius, 1793

Suborder EPIPROCTOPHORA Bechly, 1996

Subordinal clade ISOPHLEBIOPTERA Bechly, 1996

Subclade ISOPHLEBIIDA Bechly, 1996

Superfamily ISOPHLEBIOIDEA Handlirsch, 1906

Family CAMPTEROPHLEBIIDAE Handlirsch, 1920

Genus *Parasinitia* n. gen.

urn:lsid:zoobank.org:act:FD3E6C8A-A478-4FDF-80CF-ADD907B84906

**DIAGNOSIS.** — One row of cells in basal areas between MAa and MP, and between MP and CuAa; width of basal areas between MAa and MP, and between MP and CuAa nearly identical; CuAa approaching MP, with at least three rows of cells between it and posterior wing margin distally; area between MAa and MP widened before N, with at least three rows of cells in broadest area; MP with many posterior or intercalary veins, with 15 rows of cells between it and posterior wing margin in broadest area; 'O' seven cells distal of Sn; a strong and long oblique vein between IR1 and RP2, oriented towards the wing base, lying slightly close to level of Pt base than to IR1 base; Pt long and elongate, covering six cells and not braced; two or three rows of cells between RA and RP distal of Pt.

**TYPE SPECIES.** — *Parasinitia qingyunensis* n. sp.

**ETYMOLOGY.** — Prefix "para" after the similarities with the genus *Sinitia* Pritykina, 2006.

*Parasinitia qingyunensis* n. gen., n. sp.  
(Figs 2; 3)

urn:lsid:zoobank.org:act:4FB7E6D2-D358-4F2B-AA32-70FE081B578D

**DIAGNOSIS.** — The same as for the monotypic genus.

**MATERIAL.** — Holotype, NIGP163538a, b, part and counterpart of a well-preserved forewing, with posterior wing base and middle-distal wing half preserved (Fig. 2); deposited in NIGPAS.

**TYPE LOCALITY AND STRATUM.** — Qingyun Town, Yulin City, Shaanxi Province, NW China; Yanan Formation, early Middle Jurassic (c. 174–172 Ma, Aalenian; Zhang *et al.* 2021).




FIG. 1. — Geographical sketch map showing insect locality in eastern Yulin City, north Shaanxi Province, northwestern China.

**ETYMOLOGY.** — The specific name is after Qingyun Town, main locality in the area where the specimen was found.

## DESCRIPTION

Specimen NIGP163538a, b (Figs 2; 3), middle-distal wing part of wing 41.7 mm long, width at level of N 14.4 mm; distance from wing base to Arc 6.6 mm, and from N to Pt 18.8 mm. Median area free of crossveins. CuP strong, separating submedian and subdiscoidal areas at 0.7 mm basal of Arc. DC open. Subdiscoidal area broad and transverse, free of crossveins. CuA separated from MP 9.3 mm from wing base, aligned with MAB basally. Area between CuAa and MP with one row of cells basally, and as wide as postdiscoidal area, 1.9 mm wide, distinctly narrowing distally and very broad in most distal part; CuAa with 1–3 rows of cell between it and posterior wing margin in basal area. MP sigmoidally curved. MAa slightly curved, more or less parallel with MP distal of level of N; postdiscoidal area with one row of large cells basally, expanded distally with two or three cells before N, narrower distally with one row of cells below N, and two rows distally. No secondary antenodal crossveins present between C and ScP near N; three secondary antenodal crossveins present between ScP and RA distal of A×2. Fourteen postnodal crossveins present between C and RA, not aligned with nineteen postsubnodal crossveins between RA and RP1 basal of Pt. Seven antesubnodal crossveins present in distal part of area. Seven postnodal crossveins present distal of Pt. Bqr space between RP, RP2, IR2 and oblique vein 'O' very long and narrow, with one row of cells and 15 crossveins preserved. Single oblique vein 'O' present, seven cells and 5.8 mm distal of RP2 base. RP2 curved. Nodal structures well preserved; Sn oblique and well aligned with Cr and RP2. IR1 originating seven cells and 5.5 mm distal of RP2 base; IR1 weakly zigzagged basally, more or less parallel to RP1. Pt not braced, long and narrow, sclerotized, 7.7 mm long and 0.8 mm wide.




FIG. 2. — *Parasinitsia qingyunensis* n. gen., n. sp., holotype, photographs of part and counterpart (NIGP163538a, b): **A**, middle-distal part; **B**, basal part; **C**, middle part (counterpart of A); **D**, pterostigmal part. Scale bars: 2 mm.




FIG. 3. — *Parasinitsia qingyunensis* n. gen., n. sp., holotype (NIGP163538), line drawing showing forewing venation. Scale bar: 5 mm.

Area between MAa and RP3/4 widened distally, with many cells along below Pt. Area between RP3/4 and IR2 with one row of cells basal of N level, expanded distally with six rows of cells below N. Area between IR2 and RP2 with one row of cells slightly before Pt, expanded distally with 2–4 rows of cell below Pt. Area between RP2 and IR1 with one rows of cells slightly basal of Pt base, widened distally with many rows of cells. Area between IR1 and RP1 with two rows of cells before Pt, widened distally with many rows of cells. Area between RA and RP1 before Pt end, widened distally with two or three rows of cells.

## DISCUSSION

*Parasinitia qingyunensis* n. gen., n. sp. shares the typical characters of Camptero-phlebiidae: a medium size, no secondary antenodal crossveins between C and ScP, and basally opened discoidal cell in the forewing (Nel *et al.* 2009; Li *et al.* 2012a). *Parasinitia qingyunensis* n. gen., n. sp. has similar widths of the basal parts of the postdiscoidal area between MAa and MP, and that between MP and CuA, and only one row of cells between these veins in their basal parts, only shared by the following camptero-phlebiid damsel-dragonflies: *Ctenogampsophlebia* Petrulėvičius, Huang & Nel, 2011, *Honghe* Zheng, Shi, Wang, Chang, Dou, Wang & Zhang, 2017, *Karatawia* Martynov, 1925, *Melanohypsa* Pritykina, 1968, *Parazygokaratawia* Huang, Cai & Nel, 2018, *Sinitia*, and *Zygokaratawia* Nel, Huang & Lin, 2008.

*Ctenogampsophlebia* and *Honghe* have smaller wing sizes, MAa ending on MP, and fewer rows of cells between the main veins below the pterostigma, showing differences with *Parasinitia qingyunensis* n. gen., n. sp. (Petrulėvičius *et al.* 2011; Zheng *et al.* 2017). *Karatawia* has MAa zigzagged distally, one row of cells between RA and RP1 distal of the pterostigma, MP with fewer rows of cells between it and the posterior wing margin, differing from *P. qingyunensis* n. gen., n. sp. (Nel *et al.* 1993; Li *et al.* 2012b). *Melanohypsa* Pritykina, 1968, although only preserved with wing base, is similar with *P. qingyunensis* n. gen., n. sp. in sharing with an acute distal angle of the discoidal cell and two rows of cells in the anal area; it, however, differs from *P. qingyunensis* n. gen., n. sp. in having CuAa remaining parallel with MP instead of making a strong anterior curvature towards it (Pritykina 1968; Nel *et al.* 1993). *Parazygokaratawia* and *Zygokaratawia* have smaller sizes (30.8 and 35.5 mm, respectively), MAa zigzagged and weakened distally, a short pterostigma covering two or three cells, and less rows of cells between the main veins below the pterostigma, excluding any affinities with *P. qingyunensis* n. gen., n. sp. (Nel *et al.* 2008; Huang *et al.* 2018). *Sinitia* quite resembles *P. qingyunensis* n. gen., n. sp. in having CuAa moving towards MP distally, no constriction between RP2 and IR2, many rows of cell between the main veins in the distal half of wing, plus a strong and long oblique vein between IR1 and RP2, oriented towards the wing base, and lying slightly close to the level of the pterostigmal base than to IR1 base, and a long pterostigma (Pritykina 2006; Nel *et al.*

2009). *Parasinitia qingyunensis* n. gen., n. sp. differs from *Sinitia* in having the oblique vein ‘O’ seven cells distal of Sn, two or three rows of cells between RA and RP1 distal of the pterostigma, at least three rows of cells in the broad part of postdiscoidal area basad the nodus, up to eight rows of cells in the area between IR2 and RP3/4 below the pterostigma, vs only four in *Sinitia*, and c. 15 rows of cells between MP and posterior wing margin in the broadest area, vs 13 in *Sinitia* (Pritykina 2006; Nel *et al.* 2009). Some Camptero-phlebiidae genera are based on incomplete wings in which the basal part of the postdiscoidal area is not preserved, e.g. *Sarytashia* Pritykina, 1970. This genus differs from the new fossil in the areas between IR2 and RP2 and between RP2 and IR1 much narrower (Nel *et al.* 1993).

Within Camptero-phlebiidae, *Parasinitia qingyunensis* n. gen., n. sp. has many rows of cells between the main veins in the distal part (especially for the present of two or three rows of cells between RA and RP1 distal of the pterostigma), area between MAa and MP broadened in the middle part and distally constricted, many secondary intercalary veins originating from MP, ‘O’ seven cells distal of RP2 base, a strong and long oblique vein between IR1 and RP2 oriented toward the wing base, lying slightly close to the level of Pt base than to IR1 base and Pt covering many cells, all characters shared by *Parafleckium* Li, Nel, Ren & Pang, 2012 (Li *et al.* 2012a, c). *Parafleckium* however, obviously differs from *Parasinitia qingyunensis* n. gen., n. sp. in having a broad area between MP and CuAa, which is three times the size of the postdiscoidal area, MP with fifteen rows of cells between it and the posterior wing margin in the broadest part, and CuAa with only one row of cells between it and the posterior wing margin. *Junfengi yulinensis* Zheng & Zhang, 2017 is based on a male hind wing discovered from the Yanan Formation of Yulin (Zheng *et al.* 2017). Any affinities between *Junfengi* Zheng & Zhang, 2017 and *P. qingyunensis* n. gen., n. sp. can be excluded, since the former has MAa nearly parallel with MP, MP with fewer rows of cells between it and the posterior margin, the oblique vein ‘O’ three cells distal of Sn, fewer postnodal crossveins before the pterostigma, and IR2 with only one row of cells between it and RP3/4 before the level of IR1.

In conclusion, *Parasinitia qingyunensis* n. gen., n. sp. resembles the camptero-phlebiid dragonfly *Sinitia* and *Parafleckium*, but still strongly differs from these genera. We therefore justify a new genus for this unique damsel-dragonfly.

## CONCLUSIONS

A new Camptero-phlebiidae, *Parasinitia qingyunensis* n. gen., n. sp., is described from the Middle Jurassic Yanan Formation in northern Ordos Basin, northwestern China. *Parasinitia* n. gen. exhibits most similarities with *Sinitia*, but distinguished from all other camptero-phlebiid damsel-dragonflies in having MP with fifteen rows of cells between it and the posterior wing margin, two or three rows of cells between RA and RP1 distal of the pterostigma. *Sinitia* is only recorded from the Upper Jurassic in Transbaikalia. The present discovery suggests

the existence of complex relationships between the Middle and Upper Jurassic faunas of Campteropteroptera during the Jurassic, which are still unresolved. The new discovery not only adds to the diversity of Campteropteroptera, but also provides new information useful to understand the evolution of this family during early Middle Jurassic, a period earlier than the Daohugou Biota that yields abundant Campteropteroptera.

### Acknowledgements

We offer our sincere gratitude to Mr Huang Junwei for his help during the fieldtrip investigation, the two anonymous reviewers, the associated editor, Annalisa Ferretti, and the editor-in-chief, Michel Laurin, for the very useful comments on the earlier version of the manuscript. This research was supported by the Chinese Academy of Sciences (E221020002), and National Natural Science Foundation of China (42125201, 41688103).

### REFERENCES

BECHLY G. 1996. — Morphologische Untersuchungen am Flügelgädder der rezenten Libellen und deren Stammgruppenvertreter (Insecta; Pterygota; Odonata), unter besonderer Berücksichtigung der Phylogenetischen Systematik und des Grundplanes der Odonata. *Petalura* 2: 1-402.

FABRICIUS J. C. 1793. — Entomologia systematica emendata et aucta: Secundum classes, ordines, genera, species, adjectis synonymis, locis, observationibus, descriptionibus. *Christian Gottlob Proff, Hafniae* 3: 1-488. <https://doi.org/10.5962/bhl.title.125869>

FLECK G. & NEL A. 2002. — The first isophlebioid dragonfly (Odonata: Isophlebiptera: Campteropteroptera) from the Mesozoic of China. *Palaeontology* 45 (6): 1123-1136. <https://doi.org/10.1111/1475-4983.00278>

HANLIRSCH A. 1906-1908. — *Die fossilen Insekten und die Phylogenie der rezenten Formen*. Ein Handbuch für Paläontologen und Zoologen, Leipzig, V.W. Engelmann, 1430 p. [(published in parts between 1906 and 1908 as follows: i-vi, 1-160, pls 1-9 (May 1906); 161-320, pls 10-18 (June 1906); 321-480, pls 19-27 (August 1906); 481-640, pls 28-36 (October 1906); 641-800, pls 37-45 (February 1907); 801-960, pls 46-51 (June 1907); 961-1120 (November 1907); 1121-1280 (January 1908); vii-ix, 1281-1430 (July 1908)].

HANLIRSCH A. 1920. — Geschichte, Literatur, Technik, Palaeontologie, Phylogenie, Systematik, in SCHRÖDER C. (ed.), *Handbuch der Entomologie*. Vol. 3. Gustav Fisher, Jena: 117-306.

HUANG D., CAI C. & NEL A. 2018. — New damselfly dragonflies with “calopterygid”-like wing shape from the Middle Jurassic of China (Odonata: Isophlebioidea: Campteropteroptera). *Geobios* 51 (3): 181-186. <https://doi.org/10.1016/j.geobios.2018.04.003>

LI Y., NEL A., REN D. & PANG H. 2012a. — A new damselfly dragonfly from the Lower Cretaceous of China enlightens the systematics of the Isophlebioidea (Odonata: Isophlebiptera: Campteropteroptera). *Cretaceous Research* 34: 340-343. <https://doi.org/10.1016/j.cretres.2011.11.019>

LI Y., NEL A., REN D., ZHANG B. & PANG H. 2012b. — Reassessment of the Jurassic damselfly dragonfly genus *Karatawia* (Odonata: Campteropteroptera). *Zootaxa* 3417 (1): 64-68. <https://doi.org/10.11646/zootaxa.3417.1.4>

LI Y., NEL A., REN D. & PANG H. 2012c. — Redescription of the damselfly dragonfly *Parafleckium senjituense* on the basis of a more complete specimen (Odonata: Isophlebiptera: Campteropteroptera). *Zootaxa* 3597 (1): 53-56. <https://doi.org/10.11646/zootaxa.3597.1.7>

MARTYNOV A. V. 1925. — To the knowledge of fossil insects from Jurassic beds in Turkestan. *Izvestiya RAN* 17: 569-598.

NEL A., MARTINEZ-DELCLÓS X., PAICHELER J.-C. & HENROTAY M. 1993. — Les ‘Anisozygoptera’ fossiles. Phylogénie et classification (Odonata). *Martinia Numéro Hors Série* 3: 1-311.

NEL A., HUANG D. & LIN Q. 2008. — A new genus of isophlebioid damselfly dragonflies with “calopterygid”-like wing shape from the Middle Jurassic of China (Odonata: Isophlebioidea: Campteropteroptera). *European Journal of Entomology* 105: 783-787.

NEL A., BECHLY G., MARTINEZ-DELCLÓS X. & HUANG D. 2009. — New and poorly known Mesozoic damselfly dragonflies (Odonata: Isophlebioidea: Campteropteroptera, Isophlebiidae). *Palaeodiversity* 2: 209-232.

PETRULEVIČIUS J. F., HUANG D. & NEL A. 2011. — A new genus and species of damselfly dragonfly (Odonata: Isophlebioidea: Campteropteroptera) in the Middle Jurassic of Inner Mongolia, China. *Acta Geologica Sinica* 85 (4): 733-738. <https://doi.org/10.1111/j.1755-6724.2011.00478.x>

PRITYKINA L. N. 1968. — Dragonflies from Karatau (Odonata), in PANFILOV D. V. (ed.), *Fossil insects from Karatau*. Academy of Sciences of the Soviet Union, Nauka, Moscow: 1-252.

PRITYKINA L. N. 1970. — Triassic and Jurassic dragonflies of the Liassophlebiidae from Soviet Central Asia. *Paleontologicheskii Zhurnal* 4: 91-102.

PRITYKINA L. N. 2006. — Isophlebiid dragonflies from the Late Mesozoic of eastern Transbaikalia (Odonata: Isophlebiidae). *Paleontological Journal* 40: 636-645. <https://doi.org/10.1134/S0031030106060062>

RIEK E. F. 1976. — A new collection of insects from the Upper Triassic of South Africa. *Annals of the Natal Museum* 22 (3): 791-820. [https://hdl.handle.net/10520/AJA03040798\\_608](https://hdl.handle.net/10520/AJA03040798_608)

RIEK E. F. & KUKALOVÁ-PECK J. 1984. — A new interpretation of dragonfly wing venation based upon Early Carboniferous fossils from Argentina (Insecta: Odonatoidea) and basic character states in pterygote wings. *Canadian Journal of Zoology* 62 (6): 1150-1166. <https://doi.org/10.1139/z84-166>

ZHANG Z., WANG T., RAMEZANI J., LV D. & WANG C. 2021. — Climate forcing of terrestrial carbon sink during the Middle Jurassic greenhouse climate: Chronostratigraphic analysis of the Yan’an Formation, Ordos Basin, North China. *Geological Society of America Bulletin* 133 (7-8): 1723-1733. <https://doi.org/10.1130/B35765.1>

ZHENG D., NEL A., WANG B., JARZEMBOWSKI E. A., CHANG S.-C. & ZHANG H. 2016. — A new damselfly dragonfly from the Lower Jurassic of northwestern China and its palaeobiogeographic significance. *Journal of Paleontology* 90 (3): 485-490. <https://doi.org/10.1017/jpa.2016.59>

ZHENG D., DONG C., WANG H., YE Y., WANG B., CHANG S.-C. & ZHANG H. 2017. — The first damselfly dragonfly (Odonata: Isophlebioidea: Campteropteroptera) from the Middle Jurassic of Shaanxi Province, Northwestern China. *Alcheringa* 41 (4): 509-513. <https://doi.org/10.1080/03115518.2017.1318170>

ZHENG D., SHI T., WANG H., CHANG S.-C., DOU L., WANG B. & ZHANG H. 2018. — *Hongheia xui* gen. et sp. nov., the second earliest Jurassic damselfly dragonfly (Odonata: Campteropteroptera) from the Junggar Basin, NW China. *Comptes Rendus Palevol* 17 (6): 346-350. <https://doi.org/10.1016/j.crpv.2018.01.002>

ZHENG D., WANG H., NEL A., DOU L., DAI Z., WANG B. & ZHANG H. 2019. — A new damselfly dragonfly (Odonata: Anisozygoptera: Campteropteroptera) from the earliest Jurassic of the Junggar Basin, northwestern China. *Alcheringa: An Australasian Journal of Palaeontology* 43 (4): 563-567. <https://doi.org/10.1080/03115518.2019.1623321>

Submitted on 8 April 2022;  
accepted on 20 September 2022;  
published on 11 July 2023.