

Oribatid mites from deep soils of Hòn Chông limestone hills, Vietnam: the family Lohmanniidae (Acari: Oribatida), with the descriptions of *Bedoslohmanna anneae* n. gen., n. sp., and *Paulianacarus vietnamese* n. sp.

Nestor FERNANDEZ

National Council of Scientific and Technological Research (CONICET),
Subtropical Biological Institut (IBS),
Evolutive Genetic Laboratory FCEQyN, Misiones National University,
Felix de Azara 1552, 3300 Posadas Misiones (Argentina)
nestorfernand51@yahoo.fr

Pieter THERON

Research Unit for Environmental Sciences and Management,
North-West University,
2520 Potchefstroom Campus (South Africa)
pieter.theron@nwu.ac.za

Christine ROLLARD

Muséum national d'Histoire naturelle,
Département Systématique et Évolution, UMR 7205
case postale 53, 57 rue Cuvier F-75231 Paris cedex 05 (France)
chroll@mnhn.fr

Elio Rodrigo CASTILLO

National Council of Scientific and Technological Research (CONICET),
Subtropical Biological Institut (IBS),
Evolutive Genetic Laboratory FCEQyN, Misiones National University,
Felix de Azara 1552, 3300 Posadas Misiones (Argentina)
castillo.eliorodrigo@gmail.com

Fernandez N., Theron P., Rollard C. & Castillo E. R. 2014. — Oribatid mites from deep soils of Hòn Chông limestone hills, Vietnam: the family Lohmanniidae (Acari: Oribatida), with the descriptions of *Bedoslohmanna anneae* n. gen., n. sp., and *Paulianacarus vietnamese* n. sp. *Zoosystema* 36 (4): 771-787. <http://dx.doi.org/10.5252/z2014n4a5>

ABSTRACT

In this first work on the oribatid mite fauna of Hòn Chông limestone hills, Vietnam, we describe *Bedoslohmanna anneae* n. gen., n. sp. and *Paulianacarus vietnamese* n. sp., in both cases studied with optical microscopy. *Bedoslohmanna* n. gen. presents a very interesting “leg-folding process”, unusually shaped legs and significant notogastral neutrichy. *Paulianacarus* Balogh, 1960 is newly reported from Vietnam and *P. vietnamese* n. sp. is most similar to *P. rugulosus* Mahunka, 1995 from Thailand.

KEY WORDS

Acari,
Oribatida,
Lohmanniidae,
Vietnam,
new genus,
new species.

RÉSUMÉ

Les acariens Oribates des sols profonds des collines calcaires de Hòn Chông, Vietnam: la famille des Lohmanniidae (Acari: Oribatida), avec la description de Bedoslohmanna annea n. gen., n. sp., et Paulianacarus vietnamese n. sp.

Dans ce premier travail sur la faune Oribatologique des collines calcaires de Hòn Chông du Vietnam, nous décrivons *Bedoslohmanna annea* n. gen, n. sp. et *Paulianacarus vietnamese* n. sp., tous deux étudiés en microscopie optique. *Bedoslohmanna* n. gen. présente un système très intéressant pour replier ses pattes, une forme de pattes très particulière ainsi qu'une importante néotriche notogastral. *Paulianacarus* Balogh, 1960 est nouveau pour le Vietnam et *P. vietnamese* n. sp. ressemble à *P. rugulosus* Mahunka, 1995 de Thaïlande.

MOTS CLÉS

Acariens
Oribates
Lohmanniidae
Vietnam
genres nouveaux,
espèces nouvelles.

INTRODUCTION

The Hòn Chông limestone hills of Kien Giang Province, located in the extreme southwestern part of Vietnam, harbours cave and soil invertebrate communities exceptional in richness and micro-endemics (Deharveng *et al.* 2009). These tiny hills (the largest one has an area of less than one km²) are currently under intense pressure due to quarrying, and a large part of their fauna is critically threatened, with high risk of species extinctions. In spite of the large number of genera and species described from this area during the last two decades, plenty of remarkable taxa are yet to be described. The most diversified soil mites, principally oribatids, are also the least studied. A single species of mite (Actinotrichida, Eutrombidiidae) has been described so far from the area (Makol & Gabrys 2005) and another has been identified (Deharveng *et al.* 2009), whereas hundreds of species remain to be examined in the samples at hand.

Studying the biodiversity of Hòn Chông limestone hills is a very important consideration in the impact assessment of this small and fragile ecosystem.

The extreme isolation of these karst hills can generate endemism among arthropods (Deharveng *et al.* 2009) as in other isolated karsts of tropical regions.

“The number of arthropod species collected in Hòn Chông is approaching 500, most of them undescribed. Actually, this number strongly underestimates the total biodiversity of the area, because data resolution is extremely uneven from one group to the other, ranging from species or morphospecies (Collembola, Pauropoda, cave groups...) the family (many millipedes and centipedes) or even order (many spiders, mites, aquatic fauna). Mites in particular have hardly been studied though they would certainly provide the largest diversity of all soil arthropods” (Deharveng *et al.* 2009: 37).

Bedoslohmanna annea n. gen., n. sp., described in the present paper displays a very particular type of setae covering the body and legs. At the moment of deployment of the “protection mechanism” (Fernandez *et al.* 2013a), the legs are concealed behind the large umbrella-shaped setae present on all legs. The notogaster, ventral region and legs also show interesting particularities.

In *Paulianacarus vietnamese* n. sp., all setae are small and the anal and adanal plates are similar to those in *Paulianacarus rugulosus* Mahunka, 1995 from Thailand.

Due to the limited number of specimens, their rarity and particularities, we did not perform Scanning Electron Microscopy studies, which would result in the loss of the coated material, thus depositing the largest possible number of specimens in the MNHN.

MATERIAL AND METHODS

The material was obtained by Louis Deharveng and Anne Bedos using Berlese–funnel extractors (see particulars under each species) in 2003. Specimens were macerated in lactic acid, and observed in the same medium using the open-mount technique (cavity slide and cover slip) described by Grandjean (1949) and Krantz & Walter (2009). Drawings were made by means of an Olympus BHC compound microscope (Rungis, France), equipped with a drawing tube.

Measurements taken: total length (tip of rostrum to posterior edge of notogaster) and width (widest part of notogaster), in micrometres (μm). Setal formulae of the legs include the number of solenidia (in parentheses); tarsal setal formulae include the famulus (ϵ). In order to study several structures, principally notogaster, subcapitulum and legs, and particular setae, specimens were dissected and monitored during the lactic acid maceration process (in warm 70% lactic acid), before being stained with chlorazol black E (Coineau 1974).

ABBREVIATIONS

Morphology

Morphological terms and abbreviations used are those developed by Grandjean (1950); Norton & Behan-Pelletier (2009); Fernandez *et al.* (2013a–c).

Institutions

MNHN Muséum national d'Histoire naturelle, Paris.

SYSTEMATICS

Family LOHMANNIIDAE Berlese, 1916

Genus *Bedoslohmanna* n. gen.

TYPE SPECIES. — *Bedoslohmanna annea* n. sp.

ETYMOLOGY. — The generic prefix “Bedos” and the specific epithet are in homage to Dr Anne Bedos, MNHN.

DIAGNOSIS (ADULT)

Notogastral neotrichy

Genital plate undivided, ten pairs of setae; anal and adanal plate fused, with six pairs of setae, four pairs

antiaxial and two pairs paraxial situated in the middle posterior zone; preanal plate broadly triangular, posterior part covering the anterior anal-adanal plate zone; preanal plate, anterior part fused on posterior genital opening margin. Leg ventral blades: femora I and II with two prominences; femora III and IV with teeth; all femora presenting umbrella-shaped setae that conceal legs during folding process.

Bedoslohmanna annea n. sp.

(Figs 1–4; Table 1)

TYPE MATERIAL. — **Holotype:** Vietnam, Kien Giang Province; Kien Luong; Nui Hon Chong “sommet est, flanc nord”, 27.I.2003, secondary forest, 1 ♀, Berlese funnel extraction of deep mineral soil. Louis Deharveng and Anne Bedos leg. (VIET-929) X 104.647086; Y 10.140744, alt. 120 m. **Paratypes:** same data, 2 ♀♀. MNHN, preserved in 70% ethanol.

DIAGNOSIS (ADULT FEMALE)

Setation

Umbrella-shaped and simple. Umbrella-shaped: with two variations, one with rounded and more or less smooth edge, the other with spiky edge. Smooth edge: ro, le, exa, exp; notogastral, subcapitular h, m₂; epimeric; genital antiaxial; adanal; spiky edge: in setae. Simple setae: subcapitular a, m₁; paraxial genital; paraxial adanal. Umbrella-shaped setae covering almost entire body and legs.

Prodorsum

Triangular to polyhedral in dorsal view, anteriorly truncated in lateral view; postbothridial transverse band sb, poorly developed. Rostrum rounded; bothridium large, dorso-laterally opened; sensillum setiform, particularly curved, short barbs placed distally; in inserted between bothridia; Notogaster. Transverse bands not observed; setae 31–35 pairs: twelve pairs primary; 21–23 pairs additional neotrichial setae; all umbrella-shaped; c₁, c₂, c₃, d₁, d₂, cp, e₁, e₂, f₁, f₂, h₁, h₂ large; neotrichal setae small. Five pairs of lyrifissures present, ia, ip, below unsclerotized lateral longitudinal line; ips on adanal fold band; im above e₂ setae and ih posterior to h₂ setae.

Ventral region

Subcapitular setae, four pairs. Adoral setae: or_1 largest, with apical bead; or_2 elongate; or_3 small, sharply tipped. Chelicera: fixed digit, four teeth; chb seta well developed, cha seta only represented by insertion. Apodemata short, clearly visible; epimeric setal formulae 3-1-3-4. Anal setae two pairs; adanal setae four pairs.

Lateral region

Pedofossa housing retracted legs I-IV clearly visible; unsclerotized longitudinal line, terminating at level of ih lyrifissure.

Legs

All claws monodactyle, with tooth antiaxially. Femur I, II with three umbrella-shaped setae; III, IV, with one only. Setal formulae I (0-4-3-4-17-1) (2-1-2); II (0-5-3-4-14-1) (1-1-2); III (2-4-2-3-12-1) (0-1-1); IV (2-3-2-2-12-1) (1-0-0). Solenidia setiform; d setae present on tibia and genu I, II, III and only on genu IV.

DESCRIPTION

Measurements

530 μm (541–519) \times 243 μm (260–230) (Holotype 1 ♀ and 2 ♀♀ paratypes measured).

Shape

Oval (Fig. 1A, B).

Colour

Specimens without setae: yellow to light brown; slightly shiny, when observed in reflected light.

Cerotegument

Almost nonexistent; only small amount present near insertion on first and second legs.

Integument

Polyhedral microsculpture on prodorsum, zone around bo, only visible in transmitted light.

Setation (legs not included; see Legs and Folding process)

Two types: umbrella-shaped and simple setae.

Two variants of umbrella shape: one with rounded and more or less smooth edge, the second with spiky edge. Smooth-edged variant found on: prodorsum (Fig. 1B, ro, le, exa, exp setae); notogaster (Fig. 1B, all setae); subcapitular (Fig. 2E, h, m_2); epimeric (Fig. 2B, all setae); genita (Fig. 2D, (antiaxial setae); adanal setae (Fig. 2D). Spiky-edged variant: only in setae (Fig. 1B).

Simple setae on: subcapitulum, a, m_1 (Fig. 2E); paraxial zone of genital plate (Fig. 2D); paraxial setae adanal plate (Fig. 2D).

Umbrella-shaped setae presenting central zone with pedicel; upper part umbrella-shaped, with irregular margin and protuberances of different size and nerves (Figs 1B; 2B, D, E).

Prodorsum

Triangular to polyhedral in dorsal view (Fig. 1D); triangular, but anteriorly truncated in lateral view (Fig. 2A). Rostrum broadly rounded; chitinous ridge present on either side of prodorsal area (Fig. 1D). Each ro seta inserted near the anterior edge of this ridge; le setae placed near ridge, short distance posteriorly and paraxially to ro; bo large, dorsolaterally opened; si setiform, particularly curved (Figs 1A, B, D; 2A) with a few short barbs; in inserted between bothridia; exa situated near chitinous ridge. Umbrella-shaped setae covering almost entire prodorsum (Fig. 1B); shape difference between in and other prodorsal setae is notable (Fig. 1B).

Faint polygonate pattern present anterior and posterior bo. Postbothridial transverse band sb hardly discernible, only visible in specimens without setae (Fig. 1D). Dorsal view, due transparency (l.d) observed between exa and le (Fig. 1D).

Notogaster

Transverse bands not observed. Twelve pairs of primary notogastral setae: c_1 , c_2 , c_3 , d_1 , d_2 , cp, e_1 , e_2 , f_1 , f_2 , h_1 and h_2 well discernible, and between 21-23 pairs of additional neotrivial setae (Fig. 1C), situated behind h_2 . Umbrella-shaped setae covering almost entire notogaster (Fig. 1B).

Five pairs of lyrifissures present, ia, ip, situated below the unsclerotized lateral longitudinal line




FIG. 1. — *Bedoslohmannia aneae* n. gen., n. sp. adult: **A**, dorsal view, without setae; **B**, dorsal view with setae; **C**, setal neotrichy disposition; **D**, prodorsum; **E**, chelicera; **F**, fixed digit. Abbreviations: see Material and methods. Scale bars: A, B, 100 μ m; C, E, 50 μ m; D, F, 150 μ m.

(Fig. 2A) (see Lateral region); ips situated on the adanal fold band (Fig. 2C [BPAD]); im above e_2 setae and ih behind h_2 (Fig. 2C).

Lateral region

Prodorsal margin present on either side of cavities where legs I and II are housed when retracted;

TABLE 1. — *Bedoslohmanna annea* n. gen., n. sp., setae and solenidia. Abbreviations: see Material and methods.

Leg I	Femur	Genu	Tibia	Tarsus	Claw
Seta	va, vp, l', d	(l), d	d, (l), v	(p), (it), (tc), (ft), ε, pl', (pv), s, (a), (u)	1
Solenidia	-----	σ ₁ , σ ₂	-φ-	-----ω ₁ , ω ₂ -	
Leg II					
Seta	d, l', va, vp, lp	d, (l)	d, (l), v	(ft), tc', (it), (p), (u), (a), s, (pv)	1
Solenidia	-----	- σ -	-φ-	- ω ₁ , ω ₂ -	
Leg III					
Seta	(l), lp, d	l', d	d, l', v	(u), (p), (tc), (ft), a', s, (pv)	1
Solenidia	-----	- σ -	-φ-	-----0-----	
Leg IV					
Seta	lp, l', la	d, l'	d, l'	(u), (p), (tc), (it), a', s, (pv)	1
Solenidia	-----	--- σ	-0-	-----0-----	

clearly visible prodorsal posterior margin (b.p) and attachment lines of articular skin λ₁ and λ₂ (with prodorsum and notogaster respectively) (Fig. 2A).

Anterior notogastral zones presenting conspicuous tectum (Fig. 2A) and clearly defined unsclerotized lateral longitudinal line, terminating almost at level of ih lyrifissure and delimiting notaspis (unpaired dorsal) and pleuraspis (paired narrow lateral zones) (Fig. 2A); in posterior notogastral zone, when unsclerotized line does not exist, notaspis and pleuraspis not delimited (Fig. 2A).

Anteriorly each pleuraspis presenting a rounded lobe between legs II and III, where lyrifissure ia is present; posteriorly at level of d₃ and e₂ setae, well-marked edge forming roof of cavities in which legs III and IV are housed when retracted, with a protruding angle between them (Fig. 2A).

Ventral region

Subcapitulum anterior zone more or less triangular, posterior zone ovoid. Four pairs of subcapitular setae (Fig. 2E) (see Setation). Adoral setae characteristic: or₁ largest, with apical bead; or₂ elongate; or₃ with small sharp tip (Fig. 2F). Chelicera fixed digit with four teeth, one of them paraxial (indicated in Fig. 1F with dotted shading); chb setae well developed; only insertion of cha setae visible, situated laterally and posterior to chb (Fig. 1E).

Coxisternal region divided into two parts by the ventrosejugal groove. Apodemes short and clearly visible; epimeric setal formulae 3-1-3-4, all seta

of similar shape (Fig. 2B) (See Setation). Genital plate undivided with ten pairs of setae (Fig. 2D); six simple setae aligned paraxially, four umbrella-shaped setae antiaxially in semi-circle, following shape of genital plate margin. Preanal plate more or less triangular: anterior margin fused to posterior margin of genital opening, impossible to remove without breaking (Fig. 2C); posterior margin covering anterior part of ano-adanal plate, internally excavated to receive them (Fig. 2C).

Anal plate fused to adanal, resulting in single ano-adanal plate with six pairs of setae (Fig. 2C, D). Paraxial alignment of two simple setae situated in the middle posterior zone of the plate; antiaxial alignment with four umbrella-shaped setae (Fig. 2C, D). Band BPAD clearly visible in specimens immersed in lactic acid for a lengthy period; lyrifissure ips present near margin of this band (Fig. 2C, D).

Legs

Studying the legs is very complex, and due to limited number of adults available for study, it is necessary to consider our study preliminary (see Remarks).

Legs are particular by their shape, principally at femur level (Fig. 3A-D), and due to the presence of the very large umbrella-shaped setae.

Each claw bearing a single prominent tooth situated antiaxially (Fig. 3A-D).

Between tarsus and genu, all legs have a classic appearance, but femora present particular charac-




FIG. 2. — *Bedoslohmanna annea* n. gen., n. sp. adult: **A**, lateral view; **B**, ventral view (genital and ano-adanal zone, semi-schematic); **C**, genital opening, preanal plate and adanal plate without setae; **D**, genital opening, preanal plate and adanal plate with setae; **E**, subcapitulum; **F**, adoral setae. Abbreviations: see Material and methods. Scale bars: A-D, 100 μ m; E, 60 μ m; F, 20 μ m.

teristics. Two types of femur can be distinguished. Femora legs I and II displaying large ventral blade (Fig. 3A, B), with two prominences, carrying three umbrella-shaped setae. Femora legs III and IV

has ventral blade, with teeth, carrying only one umbrella-like seta (Fig. 3C, D).

Femur I robust, wide and long; ventral blade situated on proximal part, presenting two more or less




FIG. 3. — *Bedoslohmannia annea* n. gen., n. sp. adult: **A**, leg I, antiaxial; **B**, leg II, antiaxial; **C**, leg III, antiaxial; **D**, leg IV, antiaxial. Abbreviations: see Material and methods. Scale bar: 70 μ m.

rounded prominences, the distal largest; insertion of one umbrella-shaped seta on distal one. Second umbrella-like seta situated at the same level, but

more dorsally, the third towards posterior near articulation with trochanter. Femur II differing in shape; narrower and less robust, ventral blade situ-




FIG. 4. — *Bedoslohmannia anaeae* n. gen., n. sp. adult, leg folding: **A**, leg I; **B**, leg II; **C**, leg III; **D**, leg IV. Scale bar: 70 μ m.

ated further back than in femur I. Proximal prominences small, polyhedral, and carry insertion of one umbrella-shaped seta; distal prominence rounded and larger than the previous. Second umbrella-shaped seta situated distal to first, the third distal near articulation with trochanter.

Femur III presenting ventral blade with three teeth, umbrella-shaped seta situated towards distal; dorsal leaf-shaped setae present, small in relation

to umbrella-shaped setae. Femur IV displaying more or less rounded blade with only two teeth; umbrella-shaped setae situated distal.

Setal formulae I (0-4-3-4-17-1) (2-1-2); II (0-5-3-4-14-1) (1-1-2); III (2-4-2-3-12-1) (0-1-1); IV (2-3-2-2-12-1) (1-0-0).

All solenidia setiform; setae d present on tibia and genu I, II, III and genu IV, situated near solenidia (Fig. 3A-D).

Leg folding process

Depressions housing legs (l.d) are depressed zones concave-arc shaped, directing slightly obliquely, situated externally to epimeric zone (indicated by Grandjean 1950). At the moment of leg-folding, this zone partially conceals legs. Evidently the paraxial side is the unique protected zone. In this instance, depressions l.d show a complementary association with umbrella-shaped setae situated on the anti-axial surface of legs and femurs.

During leg-folding, tarsus and tibia are directed towards trochanter and femur, and the genu functions as a hinge (Fig. 4A-D). Legs are housed in depressions and large umbrella-shaped setae cover the entire anti-axial side. This protection mechanism is very simple, but effective. Umbrella-shaped epimeric setae assisting with camouflage, resulting in legs being very difficult to detect in specimens with all setae intact.

In Figure 4A-C, legs are not completely folded, permitting an understanding of positioning of legs. When fully folded, as seen in several cases, only extremities of long setae and the part of solenidia are visible (see Discussion).

REMARKS

Observation by optical microscopy is difficult due to dorsal and ventral covering of specimens by umbrella-shaped setae, which have to be removed for study.

Legs are challenging to study, because in all specimens legs were situated in leg-housing depressions, and obtaining whole legs by dissection with all setae intact was almost impossible. Preparations containing lactic acid resulted in margins of umbrella-shaped setae becoming torn. Setal shapes were similar in all cases, but sizes varied; in several cases, during observation of setae slightly from the side, they appeared to be fan-shaped structures.

Neotrichous setae are small in relation to others. Determining the number of setae affected by neotrichy necessitated removal of all, and observing insertions. A similar situation arose on the epimeric zone where all setae were removed and insertions studied.

One of the particularities of *B. anmeae* n. gen., n. sp. is the “leg-folding process” involving umbrella-shaped

setae on femora, and the leg housing depressions situated in the lateroventral zone. Processes observed in *Bovicarabodes* and *Bedoslomannia* are very different and involve dissimilar structures and strategies.

Grandjean (1950: 101) indicated that the leg housing depression, “Niches latérales pour les pattes”, permits receiving of legs when the specimen retracts them against the body. This observation by Grandjean is very important, but detailed studies on this aspect of Lohmanniidae were not made (Norton 2010 cited in page 9 and 14 this structures with the term “pedofossae”, as a characteristic of Lohmanniidae). Our initial observations created the impression that these mites had no legs as none were visible. The problem was that specimens were covered by umbrella-shaped setae and soil particles, together impeding adequate observation. Soil particles were eliminated and legs were found completely concealed behind umbrella-shaped setae on legs, and legs retracted in the leg housing depressions. Epimeric umbrella-shaped setae contributed to the protection mechanism.

On legs I and II we found three umbrella-shaped setae and on III and IV only one. In both cases, when legs are retracted to their maximum, only the apical zone of the longest setae and the solenidia are partially visible. Femoral shape and the number of umbrella-shaped setae vary among legs I-IV, possibly related to the larger number of sensorial organs to protect (in an evolutionary sense).

Genus *Paulianacarus* Balogh, 1960*Paulianacarus vietnamese* n. sp.

(Figs 5-7; Table 2)

ETYMOLOGY. — The specific epithet is dedicated to the country Vietnam, where the specimens were collected; *vietnamese* (Latin = to Vietnam).

TYPE MATERIAL. — **Holotype:** Vietnam, Kien Giang province; Kien Luong; Hon Chong, Nui Hon Chong: “nord-est de la colline, bosquet en bas de pente, terre noire fine assez sèche”, 27.I.2003, secondary forest, 1 ♀, Berlese extraction: Louis Deharveng and Anne Bedos leg. (VIET-930). X 104.649000; Y 10.139020 alt. 5 m. **Paratypes:** same data, 2 ♀♀. MNHN, preserved in 70% ethanol.




FIG. 5. — *Paulianacarus vietnamese* n. sp. adult: **A**, dorsal view; **B**, ventral view. Abbreviations: see Material and methods. Scale bar: 170 μ m.

DIAGNOSIS
 Complex microsculpture: punctillate, polyhedral. Three types of setae: *simple, small*: prodorsum, notogaster, epimeric area, genital paraxially, aggenital,

anal, adanal; *simple, large, thick*: subcapitular a, m₁, m₂; *barbate*: subcapitular h, genital antiaxially. *Prodorsum*. triangular, polyhedral; rostrum bilobate; ro setae inserted distant from rostrum; le seta posteriorly

and laterally to ro setal insertion level; bo cup-shaped, dorsally opened; si, setiform, small barbs distally; in setae posterior to bothridium and near transverse band; exa, exp positioned marginally. Postbothridial transverse band sb clearly discernible, posterior to le setae and anterior to dorsosejugal furrow.

Notogaster

Sixteen pairs of setae: $c_1, c_2, c_3, d_1, d_2, cp, e_1, e_2, f_1, f_2, h_1, h_2, h_3, p_1, p_2, p_3$. Transverse band S_2 situated posterior to c setal alignment, only band crossing sagittal plane; S_3, S_4 between c, d setal alignment; S_5 small, behind d setal alignment; S_6 curving, posterior to e_2 setal insertion; S_8 visible in ventral view, anterior to p_3 ; S_9 posterior to h_2 . Five pairs of lyrifissures: ia, ih, im, ip, ips.

Lateral region

Pleuraspis anterior zone rounded lobe; between legs I, II, zone small teeth; unsclerotized longitudinal line exceeding level ih lyrifissure.

Ventral region

Four pairs of subcapitular setae; epimeric setal formulae 3-1-3-4; genital plate undivided, 10 pairs of setae paraxially, four antiaxially; ano-adanal plate incompletely fused; anterior part of plate, presenting a very faint line, permitting identification of both plates in this zone; six pairs of setae; lyrifissure ips present as rounded structure on adanal fold band.

ADULT DESCRIPTION

Measurements

788 μm (815-790) \times 441 μm (448-432) (three specimens; deposited in MNHN). All specimens female.

Shape

Ovoid elongate (Fig. 5A, B).

Colour

Yellow to light brown; slightly shiny when observed in reflected light.

Cerotelement

Nonexistent; or disappearing during observation in lactic acid.

Integument

Complex microsculpture; punctillate cuticular surface and a little below polyhedral (Fig. 7D), microsculpture very difficult to observe on cuticular structures for example transverse band.

Setation (legs not included)

Three types: a) *simple, small*: prodorsum, notogaster, epimeric, genital paraxially, aggenital, anal, adanal, (Figs 5A, B; 6A); b) *simple, large, thick*: subcapitular a, m_1, m_2 (Fig. 5B); c) *barbate*: subcapitular h, genital antiaxially (Fig. 5B).

Prodorsum

Triangular to slightly polyhedral in dorsal view (Fig. 5A); triangular in lateral view (Fig. 6A). Rostrum with medial rounded incision, defining a bilobate rostrum (Fig. 6A; indicated by arrow), only visible in dorsal or ventral views (Figs 5A, B).

Depression housing legs l.d (for legs I and II) (Fig. 5A) well visible as concave arc-shaped zones, situated laterally (arrows indicating the course in Fig. 5); ro setae inserted far from rostrum at level of anterior arched margin of l.d (observed by transparency) (Fig. 5A); le setae placed posteriorly and laterally to ro setae insertion level; le setal insertion found at level of more arched zone of l.d (Figs 5A; 6A) (observed by transparency); bo cup-shaped, dorsally opened (Fig. 6C); si setiform, with few small barbs distally (Figs 5A; 6B); in inserted posterior to bothridium and near transverse band sb (Figs 5A; 6A); exa and exp well visible, situated marginally (Figs 5A; 6A). Transverse postbothridial band sb clearly discernible in dorsal and lateral views; situated posteriorly to le setae and anteriorly to dsj furrow; lateral end at bothridial level (Figs 5A; 6A).

Notogaster

Sixteen pairs of notogastral setae: $c_1, c_2, c_3, d_1, d_2, cp, e_1, e_2, f_1, f_2, h_1, h_2, h_3, p_1, p_2, p_3$ clearly discernible (Figs 5A; 6A).

Transverse bands: S_2 clearly visible (Figs 5A; 6A), situated posterior to c setal alignment, only band crossing sagittal plane; medial limit indistinct and variable among specimens. In dorsal view (Fig. 5A) the limit is between setal insertions c_1 and c_2 . In lateral view (Fig. 6A) the same limit almost forming




FIG. 6. — *Paulianacarus vietnamese* n. sp. adult female: **A**, lateral view; **B**, leg III antiaxial; **C**, bothridium, lateral view; **D**, genu, tibia III, dorsal view; **E**, leg IV, antiaxial. Abbreviations: see Material and methods. Scale bars: A, 350 μ m; B, E, 270 μ m; C, 23 μ m; D, 45 μ m.

TABLE 2. — *Paulianacarus vietnamese* n. sp., setae and solenidia. Abbreviations: see Material and methods.

Leg I	Femur	Genu	Tibia	Tarsus	Claw
Seta	lp, bv, v, d, l''	d, (l)	(l), x, v	(pv), m, n, s, (a), (u), (p), (tc), (ft), ε	1
Solenidia	-----	σ ₁ , σ ₂	-φ-	-----ω ₁ , ω ₂ -	
Leg II					
Seta	d, v, vp, lp, (l)	d, (l)	d, (l),	(pv), s, (a), (u), (p), (tc), (ft)	1
Solenidia	-----	-σ-	-φ-	ω	
Leg III					
Seta	d, l'	lp, la, d	v, d	ft', pl', (pv), s, (a), (u), (tc), (p)	1
Solenidia	-----	-σ-	-φ-	-----0-----	
Leg IV					
Seta	d, l', v	d, l'	d, l'	(ft), (a), pv', (tc), (p), (u), s	1
Solenidia	-----	---σ	-0-	-----0-----	

a line terminating near c_3 setal insertion; S_3 and S_4 found between c and d setal alignment (Figs 5A; 6A); S_5 small, situated posteriorly to d setal alignment and anteriorly to gla (Fig. 5); S_6 curving, situated posterior to e_2 setal insertion; S_8 only visible in ventral view (Fig. 5B) anteriorly to p_3 setal level; S_9 situated posteriorly to h_2 setal level (Figs 5A; 6A).

Five pairs of lyrifissures present: ia, ih, big size, situated below unsclerotized lateral longitudinal line; ia anteriorly on frontal lobe of pleuraspis (see Lateral region); ih situated between h_2 , f_2 setal level, well visible in lateral and ventral views (Figs 5B; 6A), in ventral view, near S_8 band, but in variable position situated either in front of or behind this band; im large, situated above unsclerotized lateral longitudinal line, near S_6 band, clearly visible in lateral view (Fig. 6A); ip small; visible in ventral view (Fig. 5B); ips rounded, situated on BPAD fold band (Fig. 5B).

Lateral region

Prodorsal margin presenting depression housing legs I,d, clearly visible (indicated with asterisks Fig. 6A) for legs I, II; bp well visible. Anterior zone of pleuraspis presenting rounded lobe with lyrifissure ia. Laterally between legs I and II, zone with small teeth (Fig. 6A, indicated with double arrow). Anterior notogastral zone presenting conspicuous tectum (Fig. 6A). Unsclerotized longitudinal line easily discernible, exceeding the level of ih lyrifissure and clearly delimiting notaspis and pleuraspis

(Fig. 6A). Situated posteriorly on same pleuraspis, l,d of legs III, IV. Posterior zone (where unsclerotized line does not exist) very small (Fig. 6A). Several muscular insertions visible (Fig. 6 indicated with black arrow).

Ventral region

Subcapitular setae, four pairs (Fig. 5B, see Setation). Apodemes well visible; epimeric setal formulae 3-1-3-4. All setae of similar shape (Fig. 2B, see Setation). Genital plate undivided with ten pairs of setae (Fig. 5B); six aligned paraxially and four anti-axially in semi-circle following shape of the genital plate margin. Preanal plate typically shaped, characteristic of the genus (Fig. 5B). Anal plate fused with adanal, delimiting single plate with six pairs of setae (Fig. 5B), anterior part of both plates with faint line, representing vestige of fusion (see Remarks). BPAD clearly visible after lengthy soaking in lactic acid; lyrifissure ips rounded, positioned near band margin (Fig. 5B).

Legs

Setal formulae I (0-5-3-4-16-1) (2-1-2); II (0-6-3-3-13-1) (1-1-1); III (2-2-3-2-13-1) (1-1-0); IV (2-3-2-2-12-1) (1-0-0).

REMARKS

Microsculpture complicating observation: limits of transverse band S_2 very hard to detect in dorsal view; in lateral view, easier, as this limit almost




FIG. 7. — *Paulianacarus vietnamese* n. sp. adult: **A**, leg I, antiaxial; **B**, palp, antiaxial; **C**, leg II, antiaxial; **D**, microsculpture. Abbreviations: see Material and methods. Scale bars: A, C, 350 μ m; B, 100 μ m; D, 2 μ m.

becomes a line. Number of genital antiaxial setae variable, on one specimen three setae on one side and four on the other.

Anal and adanal plates produce a very interesting situation: plates are fused, but anterior part with line partially delimiting anal and adanal plates; particular microsculpture of aligned points on anal plate, situated parallel, allowing differentiation of plates despite being fused, somewhat similar to

Paulianacarus rugulosus Mahunka, 1995 from Thailand. Genital setae number in *P. rugulosus* is nine, in *P. vietnamese* n. sp. asymmetric variations in the number of genital and adanal setae were observed, for example with nine-ten genital setae and four-five adanal setae. Adanal setae show variation in one specimen: one side with five and another with four.

The generic diagnosis of *Paulianacarus* was made at the same time the author instituted another new

genus, *Millotacarus* Balogh, 1960, both from Madagascar. Several authors indicated that the differences between these genera were problematic (Mahunka 1985; Coetzee 2001). Balogh & Balogh 1987 established the differences between them, adding several characters to the diagnosis of *Millotacarus*, but this did not solve the problem. The possibility exists that *Millotacarus* may be a synonym of *Paulianacarus*. The study of type material is needed.

DISCUSSION

We compared *Bedoslohmanna* n. gen. with all genera presenting the following combination of traits: undivided genital plates; fused anal and adanal plates; two pairs of anal and four pairs of adanal setae; broad preanal plates; gastronomic neotrichy and type and shape of legs.

Bedoslohmanna n. gen., showed particular characteristics: shape and disposition of preanal plate; disposition of anal and adanal setae; gastronomic neotrichy; leg shape and chaetotaxy, permitting easy differentiation of the new genus. Setal types, the leg folding process and other important characteristics observed in the new genus were not included in our comparison due to the lack of comparable studies.

The genus appearing most similar to *Bedoslohmanna* n. gen. is *Paulianacarus*, for this reason, in the present paper, species from both genera were included.

In *Bovicarabodes deharvengi* Fernandez, Theron & Rollard, 2013, *B. levyi* Fernandez, Theron & Rollard, 2013 and *B. fortidauphini* Fernandez, Theron & Rollard, 2013 (Fernandez *et al* 2013a) a very interesting “protection mechanism” was described. Several body and leg structures are involved in this process and a series of particular adaptations are observed. Evidently Carabodidae and Lohmannidae present very different characteristics and are evolutionarily distant, but in both cases a protection mechanism exists.

One of the particularities of *B. anaeae* n. gen., n. sp. is the “leg-folding process” involving umbrella-shaped setae on femora, and the leg housing depressions situated in the lateroventral zone. Processes observed in *Bovicarabodes* and *Bedoslohmanna* are

very different and involve dissimilar structures and strategies.

Grandjean (1950: 101) indicated that the leg housing depression, “Niches latérales pour les pattes” permits receiving of legs when the specimen retracts them against the body. On legs I and II we found three umbrella-shaped setae and on III and IV only one. In both cases, when legs are retracted to their maximum, only apical zone of the longest setae and the solenidia are partially visible. Femoral shape and the number of umbrella-shaped setae vary in legs I-IV, possibly related to the larger number of sensorial organs to protect (in an evolutionary sense).

Acknowledgements

This work is based on research supported in part by the National Research Foundation of South Africa (UID) 85288. Any opinion, findings and conclusions or recommendations expressed in the material are those of the authors and therefore the NRF does not accept any liability in regard thereto. The authors wish to thank the reviewers for their constructive comments.

REFERENCES

- BALOGH J. 1960. — Oribates (Acari) nouveaux de Madagascar (1^{re} série). *Mémoires de l'Institut scientifique de Madagascar*, Série A, vol. 14: 7-37.
- BALOGH J. & BALOGH P. 1987. — A new outline of the Family Lohmanniidae Berlese, 1916 (Acari, Oribatei). *Acta Zoologica hungarica* 33 (3-4): 327-398.
- COETZEE L. 2001. — New species of the family Lohmanniidae (Acari, Oribatida) from South Africa. *Navorsing van die Nasionale Museum Bloemfontein* 17 (3): 57-67.
- COINEAU Y. 1974. — *Éléments pour une monographie morphologique, écologique et biologique des Caeculidae (Acariens)*. Mémoires du Muséum national d'Histoire naturelle, 22 Zool., 81: 299.
- DEHARVENG L., BEDOS A., LE CONG KIET, LE CONG MAN & TRUONG QUANG TAM 2009. — Endemic arthropods of the Hon Chong hills (Kien Giang), an unrivalled biodiversity heritage in Southeast Asia, in LE CONG KIET, TRUONG QUANG TAM & LY NGOC SAM (eds), *Beleaguered Hills: Managing the Biodiversity of the Remaining Karst Hills of Kien Giang, Vietnam*. Nha Xuat Ban Nong Nghiep, TP. Ho Chi Minh: 31-57.
- FERNANDEZ N., THERON P. & ROLLARD C. 2013a. — The family Carabodidae (Acari: Oribatida) I. Description

- of a new genus, *Bovicarabodes* with three new species, and the redescription of *Hardybodes mirabilis* Balogh, 1970. *International Journal of Acarology* 39 (1): 26-57. <http://dx.doi.org/10.1080/01647954.2012.741144>
- FERNANDEZ N., THERON P. & ROLLARD C. 2013b. — Revision of the family Carabodidae (Acari:Oribatida) IV. *Aftibodes anjavidilavai* gen. nov., sp. nov., *Rugocephalus joffrevillei* sp. nov., and redefinition of the genus *Rugocephalus* Mahunka, 2009. *International Journal of Acarology* 39 (6): 462-480. <http://dx.doi.org/10.1080/01647954.2013.822928>
- FERNANDEZ N., THERON P. & ROLLARD C. & TIED L. 2013c. — Family Carabodidae (Acari: Oribatida) V. The genus *Congocephalus* Balogh, 1958 (second part), with a redescription of *Congocephalus involutus* Mahunka, 1997, and descriptions of two new species. *Zoosystema* 35 (4): 551-579. <http://dx.doi.org/10.5252/z2013n4a8>
- GRANDJEAN F. 1949. — Observation et conservation des très petits Arthropodes. *Bulletin du Muséum d'Histoire naturelle* 21 (2): 363-370.
- GRANDJEAN F. 1950. — Étude sur les Lohmanniidae (Oribates, Acariens). *Archives de zoologie expérimentale et générale* 87: 95-161.
- KRANTZ G. W. & WALTER D. E. 2009. — *A manual of acarology* (3rd ed). Texas Tech University Press, Lubbock, 807 p.
- MAHUNKA S. 1985. — Neue und interessante Milben aus dem Genfer Museum LIV. Oribatids from South India I (Acari: Oribatida). *Revue suisse de Zoologie* 92 (2): 367-383.
- MAHUNKA S. 1995. — New oribatids (Acari:Oribatida) from Thailand. *Acta Zoologica Academiae Scientiarum Hungaricae* 41 (2): 137-145.
- MAKOL J. & GABRYS G. 2005. — *Caecothrombium deharvengi* sp. nov. (Acari: Actinotrichida Eutrombidiidae) from Vietnam, with a proposal of Caecothrombiinae subfam. nov. *Zoologischer Anzeiger, A Journal of Comparative Zoology* (4), 243: 227-237.
- NORTON R. A. 2010. — Systematic relationships of Lohmanniidae (Acari: Oribatida), in SABELIS M. W. & BRUIN J. (eds), *Trends in Acarology*. Springer Netherlands: 175-178.
- NORTON R. & BEHAN-PELLETIER V. 2009. — Suborder Oribatida, in KRANTZ G. W. & WALTER D. E. (eds), *A manual of acarology* (3rd ed). Texas Tech University Press, Lubbock: 430-564.

*Submitted on 20 May 2014;
accepted on 31 July 2014;
published on 26 December 2014.*

