



ELSEVIER

Contents lists available at ScienceDirect

Comptes Rendus Palevol

www.sciencedirect.com



Human Palaeontology and Prehistory (Palaeohistology)

Evidence of trauma in a ca. 1-million-year-old patella of *Homo antecessor*, Gran Dolina-Atapuerca (Spain)



Preuve d'un trauma sur une rotule d'*Homo antecessor* d'environ un million d'années

Laura Martín-Francés^{a,b,*}, María Martínón-Torres^{b,c}, Ana Gracia-Téllez^{d,e}, José María Bermúdez de Castro^{a,b}

^a National Research Centre on Human Evolution (CENIEH), Paseo de la Sierra de Atapuerca 3, 09002 Burgos, Spain

^b Anthropology Department, University College London, 14 Taviston Street, WC1H 0BW London, UK

^c Universidad de Burgos, Laboratorio de Evolución Humana, Área de Paleontología, Departamento de CC. Históricas y Geografía, 09001 Burgos, Spain

^d Área de Paleontología, Departamento de Geología, Universidad de Alcalá de Henares, Edificio de Ciencias, Campus Universitario, 28871 Alcalá de Henares, Spain

^e Centro Mixto UCM-ISCIH de Investigación sobre Evolución y Comportamiento Humanos, Monforte de Lemos 5, Pabellón 14, 28029 Madrid, Spain

ARTICLE INFO

Article history:

Received 20 January 2016

Accepted after revision 10 April 2016

Available online 21 July 2016

Handled by Roberto Macchiarelli

Keywords:

Osteophyte

Trauma

Paleopathology

Patella

Homo antecessor

Mots clés :

Ostéophyte

Trauma

Paléopathologie

Rotule

Homo antecessor

ABSTRACT

We present the palaeopathological study of a left patella (ATD6-56) belonging to the Early Pleistocene species *Homo antecessor* (Atapuerca-Gran Dolina, Spain). The abnormal morphology observed in the inferior margin of the patella is an osseous overgrowth (osteophyte). Macroscopic and microscopic techniques, including microtomography and zoom stereomicroscope, were used to describe the lesion. Externally, the osteophyte has a smooth and porous texture; the boundary between the more radiolucent osteophyte and the normal bone can be identified in X-ray images. We suggest that the observed signs could be secondary to a local trauma. The lesion would have involved either the bone or related soft tissues of the left limb, possibly affecting the stability of the joint. Consequently, the individual's knee would have suffered an abnormal mechanical load that could have eventually triggered osteoarthritic changes. This is also supported by the lack of changes observed in the associated right patella (ATD6-22).

© 2016 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Nous présentons l'étude paléopathologique d'une rotule gauche (ATD6-56), appartenant à l'espèce *Homo antecessor* du Pléistocène inférieur (Atapuerca-Gran Dolina, Espagne). La morphologie anormale observée sur le bord inférieur de la rotule est une excroissance osseuse (ostéophyte). Les techniques macroscopiques et microscopiques incluant la microtomographie et le stéréomicroscope à zoom sont utilisées pour décrire la lésion. Extérieurement, l'ostéophyte a une texture douce et poreuse. La limite entre l'ostéophyte plus radiotransparent et l'os normal peut être identifiée sur les images de rayons X. Nous

* Corresponding author. National Research Centre on Human Evolution (CENIEH), Paseo de la Sierra de Atapuerca 3, 09002 Burgos, Spain.
E-mail address: lauramartinfrancesmf@gmail.com (L. Martín-Francés).

suggérons que les signaux observés pourraient être secondaires par rapport à un trauma local. La lésion aurait impliqué, soit l'os, soit les tissus tendres du membre gauche, affectant peut-être la stabilité de l'articulation. En conséquence, le genou de l'individu aurait supporté une charge mécanique anormale, qui aurait pu éventuellement déclencher des changements ostéoarthritiques. Ceci est aussi ce que corrobore l'absence de ces changements dans la rotule droite associée (ATD6-22).

© 2016 Académie des sciences. Publié par Elsevier Masson SAS. Tous droits réservés.

1. Introduction

Osteoarthritic (OA) changes including marginal osteophytes, subchondral cysts and eburnation, related to either degenerative or traumatic events (Aufderheide and Rodríguez-Martín, 1998a; Ortner, 2003a; Waldron, 2009) have been widely reported in the fossil record. The oldest examples of OA in the fossil record are those from the Pliocene *Australopithecus afarensis*, specimens AL-333w-12 and AL-333-51 (Cook et al., 1983), and *Homo ergaster*, specimens KNM-ER 803 and KNM-ER-164 (Day and Leakey, 1974). The number of OA cases documented for the Pleistocene is high. Pelvis 1 from the Middle Pleistocene sample of Atapuerca-Sima de los Huesos (SH) presents signs of OA—extensive osteophytes—in the spine. Bonmati et al. (2010) related the changes to degenerative kyphosis. Osteoarthritic changes are also frequent in Neanderthals (e.g., Dawson and Trinkaus, 1997; Gardner and Smith, 2006; Trinkaus, 1983a, b). In this species, the development of OA has been related to local factors, most likely of traumatic origin to the bone or related soft tissues. Within the Shanidar skeletal assemblage, various individuals are affected by degenerative conditions such as the postcranial skeleton of Shanidar 1, 3 and 4 (Crubézy and Trinkaus, 1992; Trinkaus, 1983a, b). La Ferrassie I was described as presenting OA changes associated with a post-traumatic episode (Dastugue and Lumley, 1976).

Probably the best-known case of extensive OA changes affecting the skeleton is that of La Chapelle-aux-Saints. Trinkaus (1985) described prominent marginal osteophytes, subchondral degeneration and eburnation in the articular facets involving numerous vertebrae. Dawson and Trinkaus (1997) suggested two scenarios for the spinal distribution of OA in this individual: either it was the consequence of an abnormal distribution of the vertebral loading, or it was related to a trauma to the lower cervical or upper thoracic vertebrae.

Gardner and Smith (2006) described numerous OA changes in several individuals from the Krapina assemblage. More recently, Trinkaus (in press) has identified and described abnormalities in eight patellar remains from Krapina, including small spines, areas of irregular subchondral bone and irregular facet surfaces. Gardner and Smith (2006) interpreted the changes observed in the Krapina elements as secondary to trauma. On the other hand, Trinkaus (in press) relate the changes observed in the patellae to the general appendicular hypertrophy of Neanderthals.

The ethio-pathological origin of an osteophyte may provide information about behavioural patterns in extinct species, such as locomotory habits or occupational

activities (e.g., Berger and Trinkaus, 1995; Lovell, 1991). For instance, the high prevalence of OA has been the basis for suggesting that Neanderthal skeletons experienced greater physical activity than other species (Trinkaus, 1978).

In this study, we report the pathological changes found in the left patella (ATD6-56) of *Homo antecessor* (Fig. 1). We provide a description, based on macroscopic and microscopic techniques, and a most likely aetiology; finally, we discuss possible behavioural implications.

2. Material and methods

2.1. Material

ATD6-56 is a left patella 37.2 mm (maximum height) × 19 mm (maximum thickness) (Carretero et al., 1999) (Fig. 1). The bone presents erosional damage of taphonomic origin on the medial margin of the articular surface; thus, the estimated breadth is 34.1 mm. The patella ATD6-56 was recovered from the TD6 level of the Atapuerca-Gran Dolina site (Burgos, Spain). The complete fossil assemblage from this site comprises more than 150 human remains assigned to *Homo antecessor*, as well as stone tools and faunal remains (Bermúdez de Castro et al., 1997; Bermúdez de Castro et al., 2015; Carbonell et al., 1995; Carretero et al., 1999). The minimum number of individuals (MNI) of the Gran Dolina-TD6 level, based on dental remains, is eight with an age range from infants to young adults (Bermúdez de Castro et al., 2015). Thermoluminescence, ESR and U-series dating methods provided a date for the site of ca. 900,000 years (Berger et al., 2008; Falguères et al., 1999).

A second patella, from the right side (specimen ATD6-22), was recovered from the same sedimentary level. Carretero et al. (1999) performed the morphological and metric analyses of both patellae (ATD6-56 and ATD6-22). Based on the morphological and metrical similarities, Carretero et al. (1999) concluded that the two small and narrow patellar bones probably belonged to the same female individual (Fig. 2). The early age of complete ossification of the patellar bone, between 14–16 years old (Scheuer and Black, 2000), prevents a more accurate age assessment. However, as the oldest individual identified so far in the TD6 sample is a young adult (Bermúdez de Castro et al., 2015), it is likely that the two patellae probably belonged to a young adult. Unfortunately, it was not possible to associate the patellae with any other lower limb remains from the site.




Fig. 1. Anterior (a), medial (b) and posterior (c) views of ATD6-56 left patella; white arrows point to the bone overgrowth (osteophyte).

Fig. 1. Vues antérieure (a), médiale (b) et postérieure (c) de la rotule gauche ATD6-56 ; les flèches blanches indiquent l'excroissance de l'os (ostéophyte).

2.2. Methods

In addition to the macroscopic analysis of the bone, we also used microscopic techniques. A zoom stereo microscope (Olympus SZ61) with camera (Olympus UC30), housed at the Microscopy Laboratory of the CENIEH, was used to visualise bone growth. In addition, ATD6-56 was scanned with a Scanco microtomographic system (mCT 80, Scanco Medical), also housed at the CENIEH, using the following parameters for the scan: 70 kV, 140 mA, and isometric voxel size = $30 \mu\text{m}^3$. The image stack was imported into Fiji[®] software package in order to obtain an X-ray image to examine the new bone formation.

3. Results

ATD6-65 has a bone-forming lesion 15 mm (medio-lateral width) \times 4.7 mm (superior-inferior height), approximately (Fig. 1). The bone spur is located along the inferior articular margin, involving greatly the medial articular surface, and extending posteriorly (Fig. 1, white arrows). The bone texture is characterized by being both smooth and porotic (Fig. 3). On the X-ray image, we identify a radiolucent area at the level where the bone growth is located, and in contrast with the radiopaque appearance of the rest of the patella. It is also possible to identify a visible boundary between the patellar margin and the new bone growth (Fig. 4, black arrows).




Fig. 2. Anterior view of right ATD6-22 (a) and left ATD6-56 (b) patellae.

Fig. 2. Vues antérieures des rotules droite ATD6-22 (a) et gauche ATD6-56 (b).




Fig. 3. Anterior view of ATD6-56 left patella and microscopic image of the bone overgrowth (osteophyte).

Fig. 3. Vue antérieure de la rotule gauche ATD6-56 et image au microscope de la croissance osseuse (ostéophyte).




Fig. 4. X-ray images of the ATD6-56 left patella; anterior (a) and inferior (b) views; the white arrows point to the osteophyte and the black arrows point to the boundary between the patellar bone and the bone overgrowth (osteophyte).

Fig. 4. Images aux rayons X de la rotule gauche ATD6-56 ; vues antérieure (a) et inférieure (b) ; les flèches blanches indiquent la limite entre l'os rotulien et l'excroissance osseuse (ostéophyte).

4. Differential diagnosis

Bone forming lesions respond to several aetiologies; for instance, fracture healing, enthesal change, tumour lesions and/or infectious processes (Aufderheide and Rodríguez-Martín, 1998b, c; Jurmain and Villotte, 2010; Ortner, 2003b, c; Ortner, 2008). We have not identified any sign related to fracture, either in the X-rays or the surface of the bone (e.g., fracture lines or cracks along the cortical bone or callous formation). Thus, we have ruled out fracture as the cause of the lesion. Since the bone growth observed in ATD6-56 is not located in any muscular insertion site (Bowden and Bowden, 2005), we can also rule out enthesal change as a likely aetiology. We also discarded tumorous conditions based on the morphology and location of the bone growth. The osteoblastic (bone forming) nature of the ATD6-56 lesion allows us to discard osteolytic (bone destroying) tumours as an explanation. Within osteoblastic tumours we also discarded osteosarcoma, because it affects the trabecular and medullary bone, and these structures are not altered in ATD6-56. We also discard osteomas, because these are characterised by a lamellar bone structure, ivory-button morphology and

they preferentially locate at the cranium (Aufderheide and Rodríguez-Martín, 1998b; Ortner, 2003c). Finally, the lack of signs related to infectious processes (e.g., cloaca and sequestrum) also allowed us to discard infection as possible cause (Aufderheide and Rodríguez-Martín, 1998c; Ortner, 2008).

5. Diagnosis

Considering the morphology of the lesion and its location at the inferior margin of the left patella (Figs. 1, 3, 4), we suggest that it corresponds to a marginal osteophyte.

Disorders of the musculo-skeletal system are characterised by bone formation (osteophyte) mainly at the joint margin and at the ligament and tendon insertions. Osteophytes are defined as a fibrocartilage-capped bony outgrowth arising in the periosteum at the margin of the articular surfaces. Osteophytes are considered a skeletal response to stress, although it is still unclear if they are a functional adaptation or a pathological process *per se*. Primarily, the development of osteophytes is strongly related to osteoarthritic changes, although other conditions such as age and trauma can trigger their formation (Menkes and Lane, 2004; Rogers et al., 1997; Van der Kraan and Van den Berg, 2007).

6. Discussion

Specimen ATD6-56 exhibits an osteophyte at its inferior margin. Considering the lack of other associated skeletal remains to this specimen except the left patella, we suggest an array of possible sequence of events to explain the development of the bone spur (osteophyte). Risk factors to account for the development of osteophytes are age, trauma (including subtypes such as fracture, dislocation, and/or deformation), and OA (e.g., Menkes and Lane, 2004; Ortner, 2003b; Rogers et al., 1997). OA is a chronic disorder of multifactorial aetiology. Two types of factors are thought to favour the development of OA, generalised and local. Generalised factors include genetic predisposition, gender, age and obesity. Local factors comprise those conditions that result in abnormal mechanical load of a joint, such as limb misalignment or ligamentous instability (Conaghan, 2002; Östör and Conaghan, 2006; Waldron, 2009; Weiss and Jurmain, 2007). The high number of knee lesions in our species has been related not only to mechanical loading but also to the wider variety of movements of the knee joint—sliding, rolling and rotation—together with the incongruent articular surfaces (Huch et al., 1997).

If ATD6-22 and ATD6-56 belong to the same individual, age cannot be the cause for the development of the osteophyte, because ATD6-22 does not present any sign of osteophytic formation. Being a monoarticular lesion, it is likely consequence of a local event that exclusively affected the left side. Occupational tasks that require kneeling and/or squatting as well as certain activities that require repetitive and high loading exercises are believed to jeopardize knee joints (e.g., Coggon et al., 2000; Conaghan, 2002; Ezzat et al., 2013; McAlindon et al., 1999; McMillan and Nichols, 2005; Rytter et al., 2009).

Within the types of OA in the knee, tibiofemoral (TF) and patellofemoral (PF) OA have been related to occupational kneeling (Ezzat et al., 2013; Hinman and Crossley, 2007; Rytter et al., 2009). Studies showed that during knee flexion the joint forces increased, the contact areas decreased, and misalignment of the patella occurred (Hinman and Crossley, 2007; Nagura et al., 2002). This will in turn contribute to the development of degenerative changes in the knee joint among people exposed to repetitive and prolonged kneeling (Ezzat et al., 2013; Hinman and Crossley, 2007; Nagura et al., 2002). The skeletal changes include the narrowing of joint space, the development of osteophytes, and subchondral sclerosis (Buckland-Wright, 2004; Ezzat et al., 2013; Hinman and Crossley, 2007; Rytter et al., 2009). Previous reports related skeletal changes to activities and postures in prehistoric populations (e.g., Trinkaus, 1975; Ubelaker, 1979). Because the lesion is asymmetrical, that affects the left but not the right patella, we could rule out kneeling as the potential cause for the development of the osteophyte.

The development of a patellar osteophyte has been also related to soft tissue injury. Two clinical cases (Harris et al., 1995; Siddiqui and Tan, 2011) have recorded the development of an osteophyte on the inferior patellar margin, similar to the one observed in ATD6-56 left patella. According to these researchers, the primary cause of the pathology observed was a soft tissue injury—eccentric muscular contraction. The posterior development of osteophytes in the patella and in the femur would have worked as a lock to prevent the patella from recurrent superior dislocation (Harris et al., 1995; Siddiqui and Tan, 2011). A third study by Cusco et al. (2009) reported similar osteophytic formation on a patella; however, the researchers concluded that the new bone formation was secondary to tibial valgus osteomy.

Because the only two available bone elements are the patellae, we believe that the most plausible scenario for the development of the osteophyte exhibited by the ATD6-56 right patella is a local trauma. That is, the osteophyte could be a consequence of a trauma (contraction or tear) of the soft tissues (ligaments and/or muscles) resulting in extra mechanical load on the knee. However, we cannot completely rule out the possibility of the osteophyte being secondary to another type of trauma. The fracture of associated limb bones—such as the left femur and/or tibia—could have also triggered a similar course of events. This interpretation remains speculative until other bone remains can be associated with this specimen. The lesion here described would be the second one related to strenuous physical activity recorded in the *H. antecessor* sample. Previously Martín-Francés et al. (2015) described a metatarsal stress fracture in specimen ATD6-124, a condition related to high-intensity, repetitive load physical activity.

7. Conclusion

The bone formation in the left ATD6-56 patella can be described as an osteophyte. We believe that the most plausible explanation for the patella involvement is a local factor, such as trauma to the left limb. The event would have affected the left but not the right side of the individual. A soft tissue lesion such as ligament tear or muscular

contraction, or the fracture of an associated bone, such as the femur or the tibia, could have added a higher mechanical load to one of the knee joints. Primary risk factors for the development of knee injuries are repetitive, high loading exercises. Moreover, recently other example of lesion related to high-intensity and repetitive load physical activities was identified in a right metatarsal of *H. antecessor* species. Despite the suggested aetiology, we are aware that more lower limb remains are needed to support the presented hypothesis. Ultimately, this report contributes to increase our knowledge about the paleopathological record, as well as aspects on the health and behaviour/habits of the Pleistocene hominins.

Funding

Grant sponsorship: Spanish Ministry of “Ciencia e Innovación”, Ministry of Culture of the “Junta de Castilla y León” (Spain), Fundación Atapuerca.

Acknowledgments

The authors would like to thank the anonymous reviewer for the comments that helped to improve the manuscript. We acknowledge all members of the Atapuerca Research Team their dedication and effort. We are grateful to Elena Lacasa-Marquina and Pilar Fernández-Colón, from the CENIEH Conservation and Restoration Department, for their invaluable work with the fossil collection. Special thanks to José Miguel Carretero for providing the image for Fig. 2. The mCT scanner of the patella was performed in the Microscopy Laboratory at CENIEH facilities with the collaboration of CENIEH staff. This research was supported with funding from the Dirección General de Investigación of the Spanish Ministerio de Educación y Ciencia (MEC), Project No. CGL2015-65387-C3-3-P, Junta de Castilla y León Projects No. BU005A09 and GR249 and The Leakey Foundation through the personal support of Gordon Getty (2013) and Dub Crook (2014, 2015 and 2016) to one of the authors (M.M.-T.). Fieldwork at Atapuerca is supported by the Consejería de Cultura y Turismo of the Junta de Castilla y León and the Fundación Atapuerca. Laura Martín-Francés has a Post-Doctoral Grant from the Fundación Atapuerca. Ana Gracia-Téllez has a Contract-Grant from the Ramón y Cajal Program, RYC-2010-06152. The authors that participated in the study and contributed to the elaboration of this manuscript declare and certify that there is no personal relationship with any financial organization or people regarding the material discussed in the manuscript.

References

- Aufderheide, A.C., Rodríguez-Martín, C., 1998a. Joint diseases. In: Aufderheide, A.C., Rodríguez-Martín, C. (Eds.), *The Cambridge Encyclopedia of Human Paleopathology*. Cambridge University Press, Cambridge, pp. 93–116.
- Aufderheide, A.C., Rodríguez-Martín, C., 1998b. Neoplastic conditions. In: Aufderheide, A.C., Rodríguez-Martín, C. (Eds.), *The Cambridge Encyclopedia of Human Paleopathology*. Cambridge University Press, Cambridge, pp. 371–392.
- Aufderheide, A.C., Rodríguez-Martín, C., 1998c. Infectious diseases. In: Aufderheide, A.C., Rodríguez-Martín, C. (Eds.), *The Cambridge*

- Encyclopedia of Human Paleopathology. Cambridge University Press, Cambridge, pp. 117–246.
- Berger, T.D., Trinkaus, E., 1995. Patterns of trauma among the Neandertals. *J. Archaeol. Sci.* 22, 841–852.
- Berger, G.W., Pérez-González, A., Carbonell, E., Arsuaga, J.L., Bermúdez de Castro, J.M., Ku, T.L., 2008. Luminescence chronology of cave sediments at the Atapuerca paleoanthropological site, Spain. *J. Hum. Evol.* 55, 300–311.
- Bermúdez de Castro, J.M., Arsuaga, J.L., Carbonell, E., Rosas, A., Martínez, I., Mosquera, M., 1997. A hominid from the Lower Pleistocene of Atapuerca, Spain: possible ancestor to Neandertals and modern humans. *Science* 276, 1392–1395.
- Bermúdez de Castro, J.M., Martínón-Torres, M., Martín-Francés, L., Modesto-Mata, M., Martínez de Pinillos, M., García, C., Carbonell, E., 2015. *Homo antecessor*: The state of the art eighteen years later. *Quatern. Int.*, <http://dx.doi.org/10.1016/j.quaint.2015.03.049>.
- Bonmatí, A., Gómez-Olivencia, A., Arsuaga, J.L., Carretero, J.M., Gracia, A., Martínez, I., Lorenzo, C., Bermúdez de Castro, J.M., Carbonell, E., 2010. Middle Pleistocene lower back and pelvis from an aged human individual from the Sima de los Huesos site. Spain. *Proc. Natl. Acad. Sci. U. S. A.* 107, 18386–18391.
- Bowden, B.S., Bowden, J.M., 2005. *An Illustrated Atlas of the Skeletal Muscles*. Morton Publishing Company, Englewood, 276 p.
- Buckland-Wright, C., 2004. Subchondral bone changes in hand and knee osteoarthritis detected by radiography. *Osteoarthr. Cartil.* 12, 10–19.
- Carbonell, E., Bermúdez de Castro, J.M., Arsuaga, J.L., Díez, J.C., Rosas, A., Cuenca-Bescós, G., Sala, R., Mosquera, M., Rodríguez, X.P., 1995. Lower Pleistocene hominids and artifacts from Atapuerca-TD6 (Spain). *Science* 269, 826–830.
- Carretero, J.M., Lorenzo, C., Arsuaga, J.L., 1999. Axial and appendicular skeleton of *Homo antecessor*. *J. Hum. Evol.* 37, 459–499.
- Coggon, D., Croft, P., Kellingray, S., Barrett, D., McLaren, M., Cooper, C., 2000. Occupational physical activities and osteoarthritis of the knee. *Arthritis Rheum.* 43, 1443–1449.
- Conaghan, P.G., 2002. Update on osteoarthritis part 1: current concepts and the relation to exercise. *Br. J. Sports Med.* 36, 330–333.
- Cook, D.C., Buikstra, J.E., DeRousseau, C.J., Johanson, D.C., 1983. Vertebral pathology in the Afar australopithecines. *Am. J. Phys. Anthropol.* 60, 83–101.
- Crubézy, E., Trinkaus, E., 1992. Shanidar 1: A case of hyperostotic disease (DISH) in the Middle Paleolithic. *Am. J. Phys. Anthropol.* 89, 411–420.
- Cusco, X., Seijas, R., Ares, O., Cugat, J., García-Balletbo, M., Cugat, R., 2009. Superior dislocation of the patella: a case report. *J. Orthop. Surg. Res.* 4, 1–3.
- Dastugue, J., Lumley, M.A., 1976. Les maladies des hommes préhistoriques. *La Préhistoire Française. Néolithique et Protohistorique II*, pp. 153–164.
- Dawson, J.E., Trinkaus, E., 1997. Vertebral osteoarthritis of the La Chapelle-aux-Saints 1 Neandertal. *J. Archaeol. Sci.* 24, 1015–1021.
- Day, M.H., Leakey, E.F., 1974. New evidence of the genus *Homo* from East Rudolf, Kenya (III). *Am. J. Phys. Anthropol.* 41, 367–380.
- Ezzat, A.M., Cibere, J., Koehoorn, M., Li, L.C., 2013. Association Between Cumulative Joint Loading From Occupational Activities and Knee Osteoarthritis. *Arthritis Care Res.* 65, 1634–1642.
- Falguères, C., Bahain, J.J., Yokoyama, Y., Arsuaga, J.L., Bermúdez de Castro, J.M., Carbonell, E., Bischoff, J.L., Dolo, J.M., 1999. Earliest humans in Europe: the age of TD6 Gran Dolina, Atapuerca. Spain. *J. Hum. Evol.* 37, 343–352.
- Gardner, J.C., Smith, F.H., 2006. The Paleopathology of the Krapina Neandertals. *Period. Biol.* 108, 471–484.
- Harris, N.J., Hay, S., Bickerstaff, D.R., 1995. Recurrent traumatic superior dislocation of the patella with interlocking osteophytes. *Knee* 2, 181–182.
- Hinman, R.S., Crossley, K.M., 2007. Patellofemoral joint osteoarthritis: an important subgroup of knee osteoarthritis. *Rheumatology* 46, 1057–1062.
- Huch, K., Kuettner, K.E., Dieppe, P., 1997. Osteoarthritis in ankle and knee joints. *Semin. Arthritis Rheum.* 26, 667–674.
- Jurmain, R., Villotte, S., 2010. Terminology. Entheses in medical literature and physical anthropology: a brief review [Online]. Document published online in 4th February following the Workshop in Musculoskeletal Stress Markers (MSM): limitations and achievements in the reconstruction of past activity patterns, University of Coimbra, July 2–3 2009, Coimbra, CIAS – Centro de Investigação em Antropologia e Saúde http://www.uc.pt/en/cia/msm/MSM_terminology3.
- Lovell, N.C., 1991. An evolutionary framework for assessing illness and injury in nonhuman primates. *Am. J. Phys. Anthropol.* 34, 117–155.
- Martín-Francés, L., Martínón-Torres, M., Gracia-Téllez, A., Bermúdez de Castro, J.M., 2015. Evidence of Stress Fracture in a *Homo antecessor* Metatarsal from Gran Dolina Site (Atapuerca, Spain). *Int. J. Osteoarchaeol.* 25, 564–573.
- McAlindon, T.E., Wilson, P.W.F., Aliabadi, P., 1999. Level of physical activity and the risk of radiographic and symptomatic knee osteoarthritis in the elderly: the Framingham study. *Am. J. Med.* 106, 151–157.
- McMillan, G., Nichols, L., 2005. Osteoarthritis and meniscus disorders of the knee as occupational diseases of miners. *Occup. Environ. Med.* 62, 567–575.
- Menkes, C.J., Lane, N.E., 2004. Are osteophytes good or bad? *Osteoarthr. Cartil.* 12, S53–S54.
- Nagura, T., Dyrby, C.O., Alexander, E.J., Andriacchi, T.P., 2002. Mechanical loads at the knee joint during deep flexion. *J. Orthop. Res.* 20, 881–886.
- Ortner, D.J., 2003a. Osteoarthritis and Diffuse Idiopathic Skeletal Hyperostosis. In: Ortner, D.J. (Ed.), *Identification of Pathological Conditions In Human Skeletal Remains*. Academic Press, San Diego, pp. 545–560.
- Ortner, D.J., 2003b. Trauma. In: Ortner, D.J. (Ed.), *Identification of Pathological Conditions In Human Skeletal Remains*. Academic Press, San Diego, pp. 119–178.
- Ortner, D.J., 2003c. Tumors and Tumors-like Lesions of Bone. In: Ortner, D.J. (Ed.), *Identification of Pathological Conditions In Human Skeletal Remains*. Academic Press, San Diego, pp. 503–544.
- Ortner, D.J., 2008. Differential diagnosis of skeletal lesion infectious disease. In: Pinhasi, R., Mays, S. (Eds.), *Advances in Human Paleopathology*. John Wiley & Sons Ltd, Chichester, pp. 191–214.
- Östör, A.J.K., Conaghan, P.G., 2006. Is there a relationship between running and osteoarthritis? *Int. Sport. Med. J.* 7, 75–84.
- Rogers, J., Shepstone, L., Dieppe, P., 1997. Bone formers: osteophyte and enthesophyte formation are positively associated. *Ann. Rheum. Dis.* 56, 85–90.
- Rytter, S., Egun, N., Jensen, L.K., Bonde, J.P., 2009. Occupational kneeling and radiographic tibiofemoral and patellofemoral osteoarthritis. *J. Occup. Med. Toxicol.* 4, 1–9.
- Siddiqui, M., Tan, M.H., 2011. Locked knee from superior dislocation of the patella—diagnosis and management of a rare injury. *Knee Surg. Sports Traumatol. Arthrosc.* 19, 671–673.
- Scheuer, L., Black, S., 2000. *Developmental Juvenile Osteology*. Elsevier, Academic Press, San Diego, pp. 587.
- Trinkaus, E., 1975. Squatting among the Neandertals: A problem in the behavioral interpretation of skeletal morphology. *J. Archaeol. Sci.* 2, 327–351.
- Trinkaus, E., 1978. Hard Times Among the Neanderthal. *J. Nat. Hist.* 87, 58–63.
- Trinkaus, E., 1983a. The Paleopathology of Shanidar Neandertals. In: Trinkaus, E. (Ed.), *The Shanidar Neanderthals*. Academic Press, New York, pp. 399–423.
- Trinkaus, E., 1983b. Neandertal postcrania and the adaptive shift to modern humans. In: Trinkaus, E. (Ed.), *The Mousterian Legacy: Human Biocultural Change in the Upper Pleistocene*. British Archaeological Reports, Oxford, pp. 165–200.
- Trinkaus, E., 1985. Pathology and posture of the La Chapelle-aux-Saints Neandertal. *Am. J. Phys. Anthropol.* 69, 345–354.
- Trinkaus, E., in press. The Krapina Human Postcranial Remains: Morphology, Morphometrics and Paleopathology. Faculty of Humanities and Social Sciences, University in Zagreb, 152 p.
- Ubelaker, D.H., 1979. Skeletal evidence for kneeling in prehistoric Ecuador. *Am. J. Phys. Anthropol.* 51, 679–685.
- Van der Kraan, P.M., Van den Berg, W.B., 2007. Osteophytes: relevance and biology. *Osteoarthr. Cartil.* 15, 237–244.
- Waldron, T., 2009. Diseases of Joints. In: Waldron, T. (Ed.), *Paleopathology*. Cambridge University Press, Cambridge, pp. 24–45.
- Weiss, E., Jurmain, R., 2007. Osteoarthritis revisited: a contemporary review of aetiology. *Int. J. Osteoarchaeol.* 17, 437–450.