

Volatile composition of *Clitocybe amoenolens*, *Tricholoma caligatum* and *Hebeloma radicosum*

Françoise FONSD^a, Sylvie RAPIOR^{b*}, Alain FRUCHIER^c,
Philippe SAVIUC^d & Jean-Marie BESSIÈRE^e

^aLaboratoire de Botanique et Mycologie,
Faculté de Pharmacie de Nancy / UMR - CNRS 7137 LIMOS,
Université Nancy 1, Faculté des Sciences et Techniques,
BP 239, 54506 Vandœuvre-lès-Nancy, France
Francoise.Fons@pharma.uhp-nancy.fr

^bLaboratoire de Botanique, Phytochimie et Mycologie / UMR - CNRS 5175 CEFE,
Faculté de Pharmacie, 15 avenue Charles-Flahault,
Université Montpellier I, BP 14491, 34093 Montpellier Cedex 5, France
srapior@ww3.pharma.univ-montp1.fr

^cLaboratoire de Chimie Organique, UMR 5076,
Ecole Nationale Supérieure de Chimie,
8 rue de l'Ecole Normale, 34296 Montpellier Cedex 5, France
fruchier@enscm.fr

^dUnité de Toxicologie Clinique et Toxicovigilance,
Centre Hospitalier Universitaire de Grenoble,
BP 217, 38043 Grenoble Cedex 9, France
psaviuc@chu-grenoble.fr

^eEcole Nationale Supérieure de Chimie,
8 rue de l'Ecole Normale, 34296 Montpellier Cedex 5, France
bessiere@enscm.fr

Abstract – The volatile extracts composition of fresh *Clitocybe amoenolens*, *Tricholoma caligatum* and *Hebeloma radicosum* were analysed by Gas Chromatography-Mass Spectrometry. Twenty-one, sixteen and twenty-three components were identified, respectively. Methyl-(*E*)-cinnamate was found in the three analysed mushrooms at various amounts. Methyl-(*E*)-cinnamate and methyl-benzoate as well as (*E*)-nerolidol and methyl-anthranilate were the key odorants of *C. amoenolens* floral odor. Combined methyl-(*E*)-cinnamate and indole derivatives should largely contribute to the complex floral odor of *T. caligatum* with a nauseous note when aged; the latter volatiles could be of chemotaxonomic interest for the genus *Tricholoma*. Various aromatic derivatives (benzaldehyde, 2-phenylethanal, 2-phenylethanol, phenylacetic acid) were responsible for the almond-like odor with a floral note of *Hebeloma radicosum*.

Basidiomycota / Benzaldehyde / Chemotaxonomy / Indole derivatives / Methyl benzoate / Methyl-(*E*)-cinnamate / (*E*)-nerolidol / Odor

* Correspondence and reprints: srapior@ww3.pharma.univ-montp1.fr Phone number: 33 (0) 467 548 083, Fax number: 33 (0) 467 411 940

Résumé – La composition de l'extrait volatil de *Clitocybe amoenolens*, *Tricholoma caligatum* et *Hebeloma radicosum* a été analysée sur matériel frais par chromatographie en phase gazeuse couplée à la spectrométrie de masse. Vingt-et-un, seize et vingt-trois composés ont été identifiés respectivement pour les trois champignons. Le (*E*)-cinnamate de méthyle a été trouvé dans chacun d'eux en quantités variables. Le (*E*)-cinnamate de méthyle, le benzoate de méthyle, le (*E*)-nérolidol et l'antranilate de méthyle sont les composés clés de l'odeur de *C. amoenolens*. L'association du (*E*)-cinnamate de méthyle et des dérivés indoliques semble contribuer largement à l'odeur florale complexe, devenant nauséuse avec l'âge, de *T. caligatum*; les dérivés indoliques pourraient présenter un intérêt chimio-taxonomique pour le genre *Tricholoma*. Différents dérivés aromatiques (benzaldehyde, 2-phényléthanal, 2-phényléthanol, acide phenylacétique) sont responsables de l'odeur d'amande amère avec une note florale d'*Hebeloma radicosum*.

Basidiomycota / Benzaldéhyde / Chimiotaxonomie / Dérivés indoliques / Benzoate de méthyle / (*E*)-cinnamate de méthyle / (*E*)-nérolidol / Odeur

INTRODUCTION

Many mushrooms are well-known to exhale characteristic odors which highly contribute to their mycological identification, i.e., fruity flavor of *Cantharellus cibarius* (Fons *et al.*, 2003), anise-like smell of *Hydnellum suaveolens* (Wood *et al.*, 1988), of *Gyrophragmium dunalii* (Rapior *et al.*, 2000c), and of *Clitocybe odora* (Rapior *et al.*, 2002), sweet-odor of *Hebeloma sacchariolens* (Wood *et al.*, 1992), candy-like odor of *Nolanea fructufragrans* (Largent *et al.*, 1990), nutty and earthy flavor of *Tuber melanosporum* (Flament *et al.*, 1990), fenugreek odor of *Lactarius helvius* (Rapior *et al.*, 2000a), musky to disagreeable scent of *Laetiporus sulphureus* (Rapior *et al.*, 2000b) and even putrid odor of *Phallus impudicus* (Borg-Karlson *et al.*, 1994).

These pleasant or unpleasant aromas are generated by mixture of many volatile components which make them usually complex in their perception by various descriptors. Besides the aroma of a number of mushroom species can evolve or even turn when aged (Flament *et al.*, 1990; Chiron & Michelot, 2005). In this study we have been interested in the three species *Clitocybe amoenolens*, *Tricholoma caligatum* and *Hebeloma radicosum* for their major floral or fruity-like aroma.

Clitocybe amoenolens Malençon is a toxic species responsible for the acromelalgic syndrome; several intoxications occurred recently in France (Maurienne valley) (Saviuc *et al.*, 2001, 2003; Moreau *et al.*, 2001; Bessard *et al.*, 2004) and in Italy (Abruzzo) (Leonardi *et al.*, 2002) after confusion with *Lepista inversa*. The odor of *C. amoenolens* has been described as a strong reminiscent smell with sweet floral, fruity note (pear) related to that of *Inocybe corydalina*, *I. piriadora* and *I. bongardii* (Malençon & Bertault, 1975; Bon, 1983, 1997; Charignon & Garcin, 1998), of *Hebeloma sacchariolens* (Bon, 1987; Contu *et al.*, 1999), of *Tricholoma caligatum* (Charignon & Garcin, 1998; Moreau, 2002). The *C. amoenolens* smell was also described as closed to that of benzoin (balsamic note), jasmine, orange blossom, iris and seringa or fruit drop odors sickly towards the end (Poumarat & Neville, 1993; Bon, 1997; Contu *et al.*, 1999; Moreau *et al.*, 2001). *C. amoenolens* taste has been described as sweet to slightly earthy or floury (Malençon & Bertault, 1975; Contu *et al.*, 1999). No volatile constituents of *C. amoenolens* have been previously identified.

Tricholoma caligatum (Viviani) Ricken (Fragrant *Tricholoma*) is better known, more widely found than *C. amoenolens* and consequently more described. The former has a dual odor reported as a pleasant spicy (cinnamon), balsamic (Peru or Tolu balsam), floral (narcissus, jasmine, pear blossom, whitethorn blossom) or fruity flavor (bergamot orange, orange, pear) followed by a nauseous note (Moser, 1983; Marchand, 1986; Riva, 1997). Its jasmine and pear to overripe pear scent is comparable to that of *I. bongardii*, *I. corydalina* and *I. piriadora* (Claus, 1978, Schmitt 1978; Mazza, 1998). According to Bessette *et al.* (1997), *T. caligatum* is a complex of several forms with a range of odor and taste. Lincoff (1998) states that varieties ground under oak in North America have pungent odor and disagreeable taste or slight odor and bitter taste whereas varieties found under conifers have fragrant cinnamon-like odor. Even, if it tastes the way it smells when fresh, *T. caligatum* is considered an edible species; the unpleasant-smelling varieties of the Fragrant *Tricholoma* are enjoyable once cooked (Fischer & Bessette, 1992). Other mycologists describe its taste as nutty, slightly or slowly bitter or not distinctive (Bessette *et al.*, 1997; Courtecuisse, 1999).

Hebeloma radicosum (Bulliard:Fries) Ricken (Rooting Fairy Cake) is a common inedible mushroom. Its odor is less complex and so better determined than that of the two previously reported mushrooms. *H. radicosum* exhales a strong pleasant odor of marzipan, cherry laurel or bitter almonds and has a bitter taste (Claus, 1978; Læssøe *et al.*, 1996; Bessette *et al.*, 1997; Courtecuisse, 1999). Despite their wide occurrence, *T. caligatum* and *H. radicosum* have been poorly investigated for secondary volatile metabolites.

In the present work, wild and fresh basidiocarps of *Clitocybe amoenolens*, *Tricholoma caligatum* and *Hebeloma radicosum* were investigated for volatile compounds by solvent extraction using Gas Chromatography-Mass Spectrometry (GC-MS) to identify the compounds responsible for their typical odors. The volatile compositions were compared and the chemotaxonomic value of some constituents was discussed therein.

MATERIALS AND METHODS

Materials and isolation of volatile extracts

Fresh and wild mushroom materials were collected in the fall of 2004 in Savoie (France) for *Clitocybe amoenolens*, and in the fall of 2003 in Languedoc Roussillon (France) for *Tricholoma caligatum* and *Hebeloma radicosum*. The fresh sporophores were brushed clean, crushed to fine particles (average particle size was 0.5 cm) and then immediately covered with diethyl ether (w/4v) to stop enzymatic activity and so maintain the freshness of the fungal specimens at the time of collection in the field. Solvent extraction was performed at room temperature in the darkness. The sample extracts were gently concentrated to a small volume (0.5 mL) under nitrogen stream.

GC-MS analysis

The organic extracts were analysed (1.0 µL) in duplicate by Gas Chromatography-Mass Spectrometry (GC-MS). GC-MS analyses were carried out using a gas chromatograph Hewlett-Packard (5890) and a mass selective detector

Hewlett-Packard (5971) with a potential of 70 eV for ionization by electron impact. Solvent extract analyses were performed using a 25 m × 0.20 μm × 0.13 μm polydimethylsiloxane BPX5 (Macherey-Nagel) fused silica capillary column. The injector and detector temperatures were 200°C and 270°C, respectively. The column was temperature programmed as follows: 50°C (2 min) to 230°C (3 C/min). The carrier gas was helium with a constant flow rate set close to 0.9 mL/min. All volatile components were identified by comparison with mass spectral library NBS (MacLafferty & Stauffer, 1989), literature spectra (Shibamoto, 1980; The Mass Spectrometry Data Centre, 1986; Jennings & National Institute of Standard and Technology, 1994; Adams, 1995) and our own data bank.

RESULTS AND DISCUSSION

The percentages of the volatile constituents and the retention indices are listed in Table 1. Chromatographic profiles of the organic extracts revealed, 21, 16 and 23 volatile components, which represented, 95.9, 98.2 and 94.5% of the volatile fraction for *Clitocybe amoenolens*, *Tricholoma caligatum* and *Hebeloma radicosum*, respectively (Table 1).

The GC-MS analysis revealed that *C. amoenolens* contained odorous aromatic derivatives, i.e., methyl-(*E*)-cinnamate (42.6%) and methyl benzoate (9.5%), and sesquiterpenes, i.e., β-barbatene (18.3%) and (*E*)-nerolidol (3.8%). For *T. caligatum*, we observed nine indole derivatives (84.4%), i.e., 2,4-dimethyl-1*H*-indole (29.4%) and 5-methoxy-2,4-dimethyl-1*H*-indole (29.5%) as well as methyl-(*E*)-cinnamate (10.5%). The *H. radicosum* extract was characterized by high amount of aromatic derivatives (52.6%), i.e., benzaldehyde (29.6%), 2-phenylethanal (9.0%), 2-phenylethanol (8.4%), β-barbatene (10.7%) and methyl-(*E*)-cinnamate (2.0%).

Methyl-(*E*)-cinnamate was found in the three analysed mushrooms at various levels; it seemed to be the key compound responsible for the pleasant aroma of *C. amoenolens* and *T. caligatum*. This cinnamic derivative is well-known in perfume and flavor industries for its fruity-balsamic scent (Arctander, 2003). It is responsible in perfume for “oriental” or sweet notes in high dilutions and strawberry/cherry-like fruity notes in lower dilutions. Arctander (2003) notices its frequent use in flavor for grape, cherry and strawberry notes. This ester was really found in strawberries (Ducruet *et al.*, 2001) and plums, blended with benzaldehyde (Ismail *et al.*, 1981); consequently, presence of methyl-(*E*)-cinnamate explains the fruity first note of *C. amoenolens* and *T. caligatum* usually compared to the note of several species of *Inocybe*. Moreover, the odorous substance of *I. corydalina* and *I. pyriodora* was also identified as methyl-(*E*)-cinnamate (Schmitt, 1978). These *Inocybe* species as well as *C. amoenolens* and *T. caligatum* exhale also a floral note (the second note of their scent) reported as corydale- and jasmine flowers-like. Methyl-(*E*)-cinnamate and methyl benzoate identified in *C. amoenolens* were commonly found in scent profile of *Corydalis cava* (Olesen & Knudsen, 1994) and jasmine (Christensen *et al.*, 1997); furthermore, it should be noted that methyl anthranilate detected in *C. amoenolens* was determined as one of the key odorants of the jasmine tea (Ito *et al.*, 2002). In addition, according to the flavoring properties published by Arctander (2003), methyl-(*E*)-cinnamate could also be responsible for the third balsamic note of *C. amoenolens* and *T. caligatum*

Table 1. Volatile composition (percentage) of fresh *Clitocybe amoenolens*, *Tricholoma caligatum* and *Hebeloma radicosum*.

Volatile compounds	RI ^a	<i>Clitocybe</i>	<i>Tricholoma</i>	<i>Hebeloma</i>
		<i>amoenolens</i>	<i>caligatum</i>	<i>radicosum</i>
		(%) ^b	(%)	(%)
Hexanal	790	0.8	–	1.8
Ethylbenzene	861	–	0.2	–
Styrene	887	–	0.1	–
Butyric acid	930	–	–	1.5
Benzaldehyde	949	0.1	1.0	29.6
2,3-Octadione	968	0.3	–	5.6
1-Octen-3-ol	974	–	–	2.6
3-Octanol	986	–	–	0.5
Limonene	1 020	0.3	–	–
2-Phenylethanal	1 032	1.0	–	9.0
Acetophenone	1 060	0.2	–	–
Linalool	1 093	0.3	–	–
Undecane	1 100	1.3	–	0.2
2-Phenylethanol	1 102	–	–	8.4
Methyl benzoate	1 130	9.5	–	–
Octanoic acid	1 160	–	–	1.6
Nonanoic acid	1 260	–	–	0.8
Benzoic acid	1 270	–	–	2.6
(<i>Z,E</i>)-2,4-Decadienal	1 279	–	–	0.3
Methyl-(<i>Z</i>)-cinnamate	1 292	0.1	0.9	–
Tridecane	1 300	0.8	–	1.1
(<i>E,E</i>)-2,4-Decadienal	1 309	–	–	3.9
1 <i>H</i> -Indole	1 320	–	–	1.9
Methyl anthranilate	1 331	0.2	–	–
Phenylacetic acid	1 445	0.8	–	3.0
Methyl-(<i>E</i>)-cinnamate	1 365	42.6	10.5	2.0
α -Barbatene	1 409	0.6	–	0.2
(<i>E</i>)-Cinnamic acid	1 430	2.5	–	–
β -Barbatene	1 440	18.3	–	10.7
2,4-Dimethyl-1 <i>H</i> -indole [7]	1 485	–	29.6	–
Pentadecane	1 500	1.8	0.2	0.5
NI ^c	1 537	–	–	3.5
(<i>E</i>)-Nerolidol	1 556	3.8	0.9	–
Myristic acid	1 578	8.1	–	4.8
NI (indole derivative)	1 639	–	0.9	–
NI (indole derivative)	1 660	–	3.0	–
4-Methoxymethyl-2-methyl-1 <i>H</i> -indole [10]	1 695	–	11.6	–
5-Methoxy-2,4-dimethyl-1 <i>H</i> -indole [8]	1 719	–	29.5	–
NI (indole derivative)	1 732	–	2.8	–
4-Hydroxymethyl-2-methyl-1 <i>H</i> -indole [9]	1 750	–	3.9	–
NI (indole derivative)	1 778	–	0.8	–
NI (M = 272)	1 798	2.5	–	–
5-Methoxy-4-methoxymethyl-2-methyl-1 <i>H</i> -indole [11]	1 800	–	2.3	–

^a Retention indices on polydimethylsiloxane BPX5 column.

^b Percentage of total ion current (TIC).

^c Not identified.

smell reported by a few descriptors. Lastly, it should be taken into account that this cinnamate derivative was previously recorded as responsible for the flavor of *Tricholoma matsutake* (Yajima *et al.*, 1981; Ahn & Lee, 1986; Wood & Fesler, 1986), a related Asian species of *T. caligatum*.

In *C. amoenolens*, methyl benzoate, the compound isolated from benzoin (Fernandez *et al.*, 2003) with heavy floral or fruity tones (Arctander, 2003) strengthened the three notes (floral, fruity and balsamic) already noticed for methyl-(*E*)-cinnamate. The floral and fruity notes of *C. amoenolens* were also enriched with (*E*)-nerolidol (apple, berry, lily, rose flavors), methyl anthranilate (grape, jasmine, orange blossom odors), linalool (floral, lemon notes) and limonene (fresh sweet citrusy smell) (Breheret *et al.*, 1997; Ito *et al.*, 2002; Jirovetz *et al.*, 2002; Arctander, 2003).

While methyl-(*E*)-cinnamate, benzaldehyde and (*E*)-nerolidol also justified the fragrant odor of *T. caligatum*, high levels of nine indole derivatives (84.4% entirely) were found in its volatile fraction. The latter compounds well-known for various disagreeable odors (Hilber, 1968; Rapior *et al.*, 1998; Arctander, 2003) could explain the second nauseous note of *T. caligatum*. Indole derivatives, i.e., indole, 2,4-dimethyl-1*H*-indole and 5-methoxy 2,4-dimethyl-1*H*-indole were previously reported from several *Tricholoma* species as *T. lascivum*, *T. inamoenum*, *T. sulfureum*, *T. sciodes* and *T. virgatum* (Hilber, 1968; Eizenhöfer *et al.*, 1990; Garlaschelli *et al.*, 1994; Pang & Sterner, 1996). Unlike Watson *et al.* (1986), Hilber (1968) and Rapior *et al.* (1998) stated that indole derivatives (indole, skatole, formyl indole) were responsible for the coal-tar or gas-like odor of the three first *Tricholoma* species reported above. It is worth to notice the complex odor of *T. lascivum* and *T. inamoenum* (as that of *T. caligatum*) with a fruity sweet note (jasmine, seringa) followed by a nauseous insecticidal or sulphurated note (Bon, 1988; Courtecuisse & Duhem, 1994; Gerhardt, 1999). Karagül-Yüceer *et al.* (2003) considered that indole and 3-methyl-1*H*-indole (skatole), described as mothball odor were key compounds for the unpleasant smell of rennet caseine.

In addition to odor, monomeric indole derivatives as bisindolic compounds also contribute to the taste of *Tricholoma* and *Collybia* species. Bisindolic derivatives are unstable pungent and non pungent compounds isolated from *T. sculpturatum*, *T. sciodes*, *T. virgatum* and *Collybia peronata* (Pang & Sterner, 1994; Sterner, 1994). Besides, in *T. lascivum*, the bitter component named lascivol release indolic derivatives after methanolysis according to Eizenhöfer *et al.* (1990) and Garlaschelli *et al.* (1994). As tentatively reported in Fig. 1, the volatile indole metabolites [7-11] listed in Table 1 for *T. caligatum* as well as lascivol could derive i) from the non-volatile intermediate components [1, 2] and then ii) from one of the intermediate components [3-6], biosynthesized in the polycetidic pathway. Hence, these unusual indole derivatives in the Higher Fungi Kingdom but reported for several *Tricholoma* species could be of chemotaxonomic interest for this genus. Aromatic derivatives, i.e., ethyl benzene and styrene were also detected from the *T. caligatum* extract by GC-MS as previously reported from headspace extract by GC-Olfactometry evaluation (Talou *et al.*, 2000); these very light volatile components should contribute to the mushroom aroma.

Concerning the organic extract of fresh *H. radicosum*, GC-MS analyses revealed minor amount of methyl-(*E*)-cinnamate and high level of odorous aromatic compounds derived from the shikimic pathway at various oxidized stages (benzaldehyde: almond odor; 2-phenylethanal: reminiscent odor of lilac and hyacinth; 2-phenylethanol: rose odor; phenyl acetic acid: honey-like note predominant at low concentration; Arctander, 2003). These results are slightly different from those reported by Rapior *et al.* (1996) from frozen material; thus, it is well




Fig. 1. Tentatively indole derivative biosynthesis from *Tricholoma* species.

known that the freezing changes considerably the volatile composition of fungal specimens as reported for *Hydnellum suaveolens* (Wood *et al.*, 1988). The fact that the chemical constituents responsible for the fungus odor change after freezing as well as during maturation (Flament *et al.*, 1990; Wu *et al.*, 2005) emphasizes the importance of recording the volatile extract at the time of collection in the field as it was carried out here for *C. amoenolens*, *T. caligatum* and *H. radicosum*.

This study highlighted for the first time the reminiscent floral odor of *Clitocybe amoenolens* due to methyl-(*E*)-cinnamate and methyl benzoate. Methyl-(*E*)-cinnamate as well as indole derivatives were reported as the dominant odorous constituents responsible for the complex odor of *Tricholoma caligatum*. The strong almond-like odor of *Hebeloma radicosum* was due to the major benzaldehyde but various aromatic derivatives with floral note (methyl-(*E*)-cinnamate, 2-phenylethanal, phenylacetic acid) enriched the chemical profile of this mushroom. Increasing requirements of the food and cosmetic industries for natural flavor and odor materials, respectively, higher fungi resources are of interest to naturally obtained aroma ingredients and in biotechnology to achieve nature-identical volatile components.

REFERENCES

- ADAMS R.P., 1995 — *Identification of essential oil components by gas chromatography-mass spectrometry*. Allured Publishing Corporation, Carol Stream, 469 p.
- ANH J.S. & LEE K.H., 1986 — Studies on the volatile aroma components of edible mushroom (*Tricholoma matsutake*) of Korea. *Han'guk Yongyang Siklyong Hakhoechi* 15: 253-257.
- ARCTANDER S., 2003 — *Perfume and Flavor Chemicals (Aroma Chemicals)*. Allured Publishing Corporation, Carol Stream Vol 2: from number 1791 to number 3102.
- BESSARD J., SAVIUC P., CHANE-YENE Y., MONNET S. & BESSARD G., 2004 — Mass spectrometric determination of acromelic acid A from a new poisonous mushroom: *Clitocybe amoenolens*. *Journal of Chromatography A* 1055: 99-107.
- BESSETTE A.E., BESSETTE A.R. & FISCHER D.W., 1997 — *Mushrooms of Northeastern North America*. University Press, Syracuse, NY, 582 p.
- BON M., 1983 — *Tricholomataceae* de France et d'Europe occidentale. 6^e partie : Tribu *Clitocybeae* Fay. Clé monographique. *Documents Mycologiques* 13(51): 1-53.
- BON M., 1987 — Quelques espèces intéressantes étudiées au stage FMDS de Saint-Germain-Mont-d'Or. *Bulletin de la Fédération Mycologique du Dauphiné-Savoie* 105: 28.
- BON M., 1988 — *Champignons de France et d'Europe Occidentale*. Arthaud, Paris, 368 p.
- BON M., 1997 — Flore mycologique d'Europe. Les *Clitocybes*, *Omphales* et ressemblants. *Tricholomataceae (Clitocyboideae)*. *Documents Mycologiques, Mémoire Hors Série n° 4* : 43.
- BORG-KARLSON A.K., ENGLUND F.O. & UNELIUS C.R., 1994 — Dimethyl oligosulphides, major volatiles released from *Sauromatum guttatum* and *Phallus impudicus*. *Phytochemistry* 35: 321-323.
- BREHERET S., TALOU T., RAPIOR S. & BESSIÈRE J.M., 1997 — Monoterpenes in the aromas of fresh wild mushrooms (Basidiomycetes). *Journal of Agricultural and Food Chemistry* 45: 831-836.
- CHARIGNON Y. & GARCIN R., 1998 — Un nouveau champignon toxique en France. *Bulletin de la Fédération Mycologique du Dauphiné Savoie* 149 : 11-14.
- CHIRON N. & MICHELOT D., 2005 — Odeurs de champignons : Chimie et rôle dans les interactions biotiques – une revue. *Cryptogamie, Mycologie* 26(4): 299-364.

- CHRISTENSEN L.P., JAKOBSEN H.B., KRISTIANSEN K. & MALLER J., 1997 — Volatiles emitted from flowers of γ -radiated and non radiated *Jasminum polyanthum* Franch. *in situ*. *Journal of Agricultural and Food Chemistry* 45: 2199-2203.
- CLAUS G., 1978 — Des odeurs en Mycologie. *Documents Mycologiques* 8 (30-31): 31-63.
- CONTU M., SIGNORELLO P. & ANASTASE A., 1999 — *Clitocybe amoenolens* Mal. in Abruzzo, con osservazioni sulla sua posizione sistematica. *Bollettino dell'Associazione Micologica ed Ecologica Romana* 48(3): 16-18.
- COURTECUISSÉ R., 1999 — *Mushrooms of Britain and Europe*. Harper Collins Publishers, London, 904 p.
- COURTECUISSÉ R. & DUHEM B., 1994 — *Guide des Champignons de France et d'Europe*. Delachaux et Niestlé, Paris, 476 p.
- DUCRUET V., FOURNIER N., SAILLARD P., FEIGENBAUM A. & GUICHARD E., 2001 — Influence of packaging on the aroma stability of strawberry syrup during shelf life. *Journal of Agricultural and Food Chemistry* 49: 2290-2297.
- EIZENHÖFER T., FUGMANN B., SHELDRIK W.S., STEFFAN B. & STEGLICH W., 1990 — Lascivol, der Bitterstoff des Unverschämten Ritterlings, *Tricholoma lascivum* (Agaricales). *Liebigs Annalen der Chemie* 11: 1115-1118.
- FERNANDEZ X., LIZZANI-CUVELIER L., LOISEAU A.M., PÉRICHET C. & DELBECQUE C., 2003 — Volatile constituents of benzoin gums: Siam and Sumatra. Part 1. *Flavour and Fragrance Journal* 18: 328-333.
- FISCHER D.W. & BESSETTE A.E., 1992 — *Edible wild mushrooms of North America*. University of Texas Press, Austin, USA, 254 p.
- FLAMENT I., CHEVALLIER C. & DEBONNEVILLE C., 1990 — Analysis of volatile flavor constituents of Perigord Black Truffle (*Tuber melanosporum*). *Rivista Italiana EPPOS* 9: 280-299.
- FONS F., RAPIOR S., EYSSARTIER G. & BESSIÈRE J.M., 2003 — Les substances volatiles dans les genres *Cantharellus*, *Craterellus* et *Hydnum*. *Cryptogamie Mycologie* 24: 367-376.
- GARLASCHELLI L., PANG Z., STERNER O. & VIDARI G., 1994 — New indole derivatives from the fruit bodies of *Tricholoma sciodes* and *T. virgatum*. *Tetrahedron* 50: 3571-3574.
- GERHARDT E., 1999 — *Guide Vigot des Champignons*. Editions Vigot, Paris, 714 p.
- HILBER O., 1968 — Indol als hauptkomponente des geruches einiger *Tricholoma*-arten und von *Lepiota bucknallii*. *Zeitschrift für Pilzkunde* 34: 153-158.
- ISMAIL H.M.M., WILLIAMS A.A. & TUKNOTT O.G., 1981 — The flavour components of plums: an examination of the aroma components present in the headspace above four cultivars of intact plums, Marjorie's seedling, Merton Gem, NA 10 and Victoria. *Journal of the science of food and agriculture* 32: 498-502.
- ITO Y., SUGIMOTO A., KAKUDA T. & KUBOTA K., 2002 — Identification of potent odorants in chinese Jasmine green tea scented with flowers of *Jasminum sambac*. *Journal of Agricultural and Food Chemistry* 50: 4878-4884.
- JENNINGS W. & SHIBAMOTO T., 1980 — *Qualitative analysis of flavor and fragrance volatiles by glass capillar gas chromatography*. Academic Press, New York, 472 p.
- JIROVETZ L., BUCHBAUER G., NGASSOUM M.B. & GEISSLER M., 2002 — Aroma compound analysis of *Piper nigrum* and *Piper guineense* essential oils from Cameroon using solid-phase microextraction-gas chromatography, solid-phase microextraction-gas chromatography-mass-spectrometry and olfactometry. *Journal of Chromatography A* 976: 265-275.
- KARAGÜL-YÜCEER Y., VLAHOVICH K.N., DRAKE M. & CADWALLADER K.R., 2003 — Characteristic aroma components of rennet casein. *Journal of Agricultural and Food Chemistry* 51: 6797-6801.
- LÆSSØE T., LINCOFF G. & DEL CONTE A., 1996 — *The Knopf Mushroom Book*. Alfred A. Knopf, Toronto, 256 p.

- LARGENT D.L., BRADSHAW D.E. & WOOD W.F., 1990 — The candy-like odor of *Nolanea fructufragrans*. *Mycologia* 82: 786-787.
- LEONARDI M., CIULLI G., PACIONI G. & RECCHIA G., 2002 — Una intossicazione collettiva da *Clitocybe amoenolens* riconducibile alla sindrome acromelalgica. *Micologia e Vegetazione Mediterranea* 17:133-142.
- LINCOFF G.H., 1998 — *National Audubon Society Field Guide to North American Mushrooms*. Alfred A. Knopf, New York, 926 p.
- MACLAFFERTY F.W. & STAUFFER D.B., 1989 — *The Wiley NBS Registry of mass spectral data*. Wiley & Sons, New York, 256 p.
- MALENÇON G. & BERTAULT R., 1975 — *Flore des champignons supérieurs du Maroc. Essai descriptif et critique*. Tome II (Agaricales). Editions marocaine et internationales, Tanger, Maroc, 138-141.
- MARCHAND A., 1986 — *Champignons du Nord et du Midi. Les Tricholomes*. Tome 9. Société Mycologique des Pyrénées méditerranéennes, Perpignan, France, p. 895.
- MAZZA R., 1998 — Introduzione alla « micosmologia ». *Pagine di micologia* (10): 29-96.
- MOREAU P.A., COURTECUISSE R., GUEZ D., GARCIN R., NEVILLE P., SAVIUC P. & SEIGLE-MURANDI F., 2001 — Analyse taxinomique d'une espèce toxique : *Clitocybe amoenolens* Malençon. *Cryptogamie Mycologie* 22: 95-117.
- MOREAU P.A., 2002 — Les odeurs des champignons : un monde à découvrir. *La Garance voyageuse* (61) : 33-41.
- MOSER M., 1983 — *Keys to Agarics and Boleti (Polyporales, Boletales, Agaricales, Russulales)*. Roger Phillips, London, 535 p.
- NATIONAL INSTITUTE OF STANDARD AND TECHNOLOGY, 1994 — *PC version of the NIST / EPA / NIH Mass Spectra Database*, version 4.5, U.S. Department of Commerce, Gaithersburg, Maryland.
- OLESEN J.M. & KNUDSEN J.T., 1994 — Scent profiles of flower colour morphs of *Corydalis cava* (Fumariaceae) in relation to foraging behaviour of Bumblebee queens (*Bombus terrestris*). *Biochemical and Systematic Ecology* 22: 231-237.
- PANG Z. & STERNER O., 1994 — The isolation of 2,2'-biindoline-3,3'-diones from injured fruit bodies of *Collybia peronata* and *Tricholoma scalpturatum*. *Journal of Natural Product* 57: 852-857.
- PANG Z. & STERNER O., 1996 — Novel indole derivatives from the fruit bodies of *Tricholoma sciodes*. *Acta Chemica Scandinavica* 50: 303-304.
- POUMARAT S. & NEVILLE P., 1993 — Espèce de la zone de *Quercus ilex* au Maroc, montagnarde en France : *Clitocybe amoenolens* Malençon. *Bulletin de la Fédération des Associations Mycologiques Méditerranéennes* (4): v16-19.
- RAPIOR S., BREHERET S., TALOU T., PÉLISSIER Y. & BESSIÈRE J.M., 2002 — The anise-like odor of *Clitocybe odora*, *Lentinellus cochleatus* and *Agaricus essetei*. *Mycologia* 94: 373-376.
- RAPIOR S., BREHERET S., TALOU T., PÉLISSIER Y., MILHAU M. & BESSIÈRE J.M., 1998 — Volatile components of fresh *Agrocybe aegerita* and *Tricholoma sulfureum*. *Cryptogamie Mycologie* 19: 15-23.
- RAPIOR S., CAVALIÉ S., CROZE P., ANDARY C., PÉLISSIER Y. & BESSIÈRE J.M., 1996 — Volatile components of ten frozen mushrooms (Basidiomycetes). *Journal of Essential Oil Research* 8: 63-66.
- RAPIOR S., FONTS F. & BESSIÈRE J.M., 2000a — The fenugreek odor of *Lactarius helvus*. *Mycologia* 92: 305-309.
- RAPIOR S., KONSKA G., GUILLOT J., ANDARY C. & BESSIÈRE J.M., 2000b — Volatile composition of *Laetiporus sulphureus*. *Cryptogamie Mycologie* 21: 67-72.
- RAPIOR S., MAURUC M.J., GUINBERTEAU J., MASSON C.L. & BESSIÈRE J.M., 2000c — Volatile composition of *Gyrophragmium dunalii*. *Mycologia* 92: 1043-1046.
- RIVA A., 1997 — *Tricholoma caligatum* (Viviani) Ricken 1915. Un esempio intrigante. *Bollettino dell' Associazione Micologica ed Ecologica Romana* 41-42 (2-3): 10-16.

- SAVIUC P.F., DANIEL V.C., MOREAU P.A., GUEZ D.R., CLAUSTRE A.M., CARPENTIER P.H., MALLARET M.P. & DUCLUZEAU R., 2001 — Erythromelalgia and mushroom poisoning. *Clinical Toxicology* 39: 403-407.
- SAVIUC P., FLESCHE F. & DANIEL V., 2003 — Intoxications par les champignons : syndromes majeurs. *Encyclopédie Médico-Chirurgicale. Toxicologie Pathologie Professionnelle*. Elsevier, Paris, 16-077-A-10. 10 p.
- SCHMITT J.A., 1978 — Über den Duftstoff zweier Rißpilzarten (Agaricales, Basidiomycetes). *Zeitschrift für Naturforschung* 33C: 817-819.
- STERNER O., 1994 — The isolation and structure determination of sciadole, a new indole derivative from the fruit bodies of *Tricholoma sciodes*. *Natural Product Letters* 4: 9-14.
- TALOU T., BREHERET/HULIN-BERTAUD S. & GASET A., 2000 — Identification of the major key flavour compounds in odorous wild mushrooms. *9th Weurman Flavour Research Symposium (Frontiers of Flavour Science)*. Juin 1999, Freising: 46-50.
- THE MASS SPECTROMETRY DATA CENTRE., 1986 — *Eight peak index of mass spectra*, 3rd ed. The Royal Society of Chemistry, Nottingham, 1338 p.
- WATSON R.L., LARGENT D.L. & WOOD W.F., 1986 — The “coal tar” odor of *Tricholoma inamoenum*. *Mycologia* 78: 965-966.
- WOOD W.F., BROWNSON M., SMUDDE R.A. & LARGENT D.L., 1992 — 2-Aminobenzaldehyde: the source of the “sweet odor” of *Hebeloma sacchariolens*. *Mycologia* 84: 935-936.
- WOOD W.F., DESHAZER D.A. & LARGENT D.L., 1988 — The identity and metabolic fate of volatiles responsible for the odor of *Hydnellum suaveolens*. *Mycologia* 80: 252-255.
- WOOD W.F. & FESLER M., 1986 — Mushroom odors. Student synthesis of the odoriferous compounds of the Matsutake Mushroom. *Journal of Chemical Education* 63: 92.
- WU S., ZORN H., KRINGS U. & BERGER R.G., 2005 — Characteristic volatiles from young and aged fruiting bodies of wild *Polyporus sulfureus* (Bull.: Fr.) Fr. *Journal of Agricultural and Food Chemistry* 53: 4524-4528.
- YAJIMA I., YANAI T., NAKAMURA M., SAKAKIBARA H. & HAYASHI K., 1981 — Volatile flavor compounds of Matsutake - *Tricholoma matsutake* (Ito & Imai) Sing. *Agricultural and Biological Chemistry* 45: 373-377.