

## ***Cantharellus fistulosus* sp. nov. from Tanzania**

Donatha TIBUHWA<sup>a</sup>, Bart BUYCK<sup>b\*</sup>, Amelia KIVAISI<sup>a</sup> & Leif TIBELL<sup>c</sup>

<sup>a</sup>University of Dar es salaam, Department of Molecular Biology and Biotechnology  
P.O. Box 35179, Dar es Salaam, Tanzania

<sup>b</sup>Muséum National d'Histoire Naturelle,  
Département Systématique et Evolution USM Taxonomie et Collections (CP 39),  
57, rue Cuvier, 75231 Paris Cedex 05, France

<sup>c</sup>Uppsala University, Department of Systematic Botany (EBC);  
Norbyvägen 18 D, S- 75236, Uppsala, Sweden

**Abstract** – *Cantharellus fistulosus* sp. nov. from Tanzania is described as a close relative to *C. schmitzii* from tropical miombo woodland dominated by tree species in *Brachystegia*, *Julbernardia* and *Isoberlinia*. The species is characterised by having a hollow, smooth stipe and a pink hymenium that contrasts with the yellowish brown stipe and cap surface.

***Cantharellus* / Taxonomy / Tanzania / Miombo woodland**

### INTRODUCTION

The genus *Cantharellus* Fries was described in 1821 in *Systema Mycologicum* (Fries, 1821-1832). It has been recorded from all parts of the world with the exception of the Antarctic (Danell, 1994). In Tanzania, information on occurrence and diversity of *Cantharellus* is limited to Härkönen *et al.* (1995, 2003), who reported on six species, and Buyck *et al.* (2000) who reported on 11 species and introduced two new taxa: *Cantharellus tomentosus* Eyssart. & Buyck and *C. isabellinus* var. *parvisporus* Eyssart. & Buyck.

*Cantharellus* forms ectomycorrhiza and its distribution coincides with the distribution patterns of potential mycorrhizal partners. So far, all the partners for *Cantharellus* in Tanzania are dominant trees of the Miombo woodlands, which are divided into a South-eastern and a North-western zone in the country (Fig. 1). Miombo covers an estimated area of 2.7 million km<sup>2</sup> on nutrient-poor soils in sub-Saharan Africa that receives less than 700 mm of precipitation per year (Campbell *et al.*, 1996). It is distinguished from other African savannah woodland and forest formations by the high frequency of tree species with meso- and microphyllous compound leaves (Van der Meulen and Werger, 1984), the flush of new leaves before new rains and the dominance of ectomycorrhizal tree species (Högberg, 1982, 1992; Högberg and Pearce, 1986). 'Miombo' refers to a Bantu word for *Brachystegia*, the predominant tree genus in this type of vegetation. The woodland is further characterized by the local codominance of ectomycorrhizal trees in

---

\* Corresponding author: buyck@mnhn.fr




Fig. 1. Map showing the distribution of Miombo-woodland in Tanzania. Approximate positions of collecting sites are marked with 'X'.


other genera of Caesalpinaceae, especially *Julbernardia* and *Isoberlinia*, as well as by host trees of the genus *Uapaca* (Phyllantaceae).

*Cantharellus* species not only are important for the ecology of this woodland by supplying mycorrhizal partners for trees (Munyanziza, 1994), but are also economically vital for the low income communities in the area. Various *Cantharellus* species are sold in local markets and along roadsides during the rainy season not only in Tanzania but also in the neighbouring countries (Morris, 1984; Buyck and Nzigidahera, 1995).

The objective of the present study is to describe and illustrate the small and rare *Cantharellus fistulosus* sp. nov. collected from the Miombo woodlands of Tanzania.

## MATERIALS AND METHODS

The specimens were recently collected from Miombo woodlands located in the Kazimzumbwi forest reserve Kisarawe, in the Coast region, and in the Madibira-Iringa region (Fig.1). They are found growing in small troops emerging from the forest floor, on decayed woody humus deposits beneath living trees overall dominated by the ectomycorrhizal canopy of *Brachystegia*, *Combretum* and *Julbernardia*.



Figs 2-6. *Cantharellus fistulosus* (holotype). 2. Basidia. 3. Basidiola. 4. Spores. 5. Hyphal terminations of pileus surface. 6. Fragments of hyphae just below the subhymenium with minute parietal incrustations indicated in part of one element as seen in Congo red. Scale bar = 5  $\mu$ m for spores, 10  $\mu$ m for the other elements. (all drawings B.Buyck)



Figs 7-8. *Cantharellus fistulosus*. Field aspect of fresh collections in situ. 7. Holotype. 8. Tibuhwa nr D.59 (all pictures D.Tibuhwa)

Microscopic characters were described from fresh specimens preserved in CTAB and observed in 10% ammonium solution in an aqueous solution of Congo red. Twenty measurements of basidiospores and basidia were analysed statistically to give (min) min-SD - AV -max-SD (max) Q; in which min = lowest value recorded for the measured specimen, max = highest value, AV = arithmetic average and SD standard deviation; Q the ratio length/width.

## RESULTS

*Cantharellus fistulosus* Tibuhwa & Buyck sp nov.

*Cantharello schmitzii* differt hymenophoro roseo tincto sporis haud subglobosis pileipellis hyphis valde angustioribus. Holotypus: Tanzania. Kazimzumbwi forest reserve, Kisarawe, Coast region, S 06°04'324" E 039°15'567", April 2007 on decayed woody humus in miombo woodland, Tibuhwa D 43.2007 (holotypus UPS; isotypi PC and UDSM).

**Vernacular names:** Zaramo dialect: KIZOGORO MDOGO, Hehe dialect: WISOGORO MDOGO (Literally means small *Cantharellus*)

**Cap** 1.5-2.5 cm diam., bell-shaped with strongly incurved margin when young, then expanding with age to plano-convex, becoming irregularly and slightly fissured; cap center slightly depressed and brown matted, contrasting with the bright yellow colour of the more marginal part of the cap. **Context** yellow, soft, watery and unchanging. **Hymenophore** of well-developed gills, these thick, decurrent, widely spaced, forked, pink-coloured and strongly contrasting against the yellow stipe and cap margin. **Stipe** 1.5-2.8 ¥ 0.2-0.5 cm, smooth, cylindrical, sometimes laterally compressed, with the base slightly enlarged, hollow. **Taste** slightly bitter when raw, but good when cooked. **Odour** not particular. **Spore deposit** pale pink.

**Basidiospores** ellipsoid or slightly reniform, 5-6.3-7 ¥ 3.5-3.75-4 µm, Q=1.43-1.69-2.0, smooth. **Basidia** (38)45-60(70) ¥ 5-7 µm, pedicellate-clavate, irregularly sinuous, mostly 5-6-spored. **Hymenial cystidia** not observed. **Pileipellis** composed of septate hypha measuring 4-6(7) µm diam., the terminal cells (13)25-50(65) µm long, mostly 3-5 µm diam., often narrowing slightly upward and somewhat sinuous or with a slight subapical constriction, in lower layers most hyphae up to 7 µm diam., with frequent anastomosing 'bridges' between adjacent cells of different hyphae; cell walls not exactly 'thin' but all more or less slightly thickened and therefore clearly refringent under the microscope; carotene pigment distinctly present. Near the subhymenium most hyphae again narrower, 2-4 µm diam., some with weak, yet distinct, zebroid transverse incrustations **Clamps** present everywhere.

**Habitat:** Gregarious to imbricate in small troops emerging from the *Brachystegia-Combretum-Julbernardia* woodland soil, on decayed woody humus deposits beneath the trunks of living trees, April-May.

**Specimens examined:** Tanzania, Iringa region, Madibira forest S 08°15'086" E 35°17'219" at an altitude of 1847m, in Uapaca woodland, May 2007, Tibuhwa D 59.2007 (UPS, UDSM).; Coast region, Kazimzumbwi forest reserve, Kisarawe, S 06°04'324" E 039°15'567". April 2007, miombo dominated by *Brachystegia, Combretum* and *Julbernardia*, Tibuhwa D 43.2007 UPS (holotype, isotypes: PC,UDSM).

## DISCUSSION

The species is easily recognized in the field by its small size, yellow color with clearly brown matted center, pink hymenophore composed of widely spaced gills, and by the smooth hollow stipe, which is slightly twisted or compressed.

Although too small and too rare to represent some culinary interest, the bitter taste disappears when the mushrooms are prepared.

The hollow stipe and marked 'fibrils' on the cap are reminiscent of the genus *Craterellus* Pers., but occur also in some of the smaller chanterelles of subgenus *Parvocantharellus* Eyssartier & Buyck.

Our species is morphologically very close to *C. schmitzii* Heinem. described from a very similar habitat (humus-rich trunk base) from neighboring miombo in the south of the Democratic Republic of Congo (Heinemann, 1966). The latter species – only known from the type collection – differs in particular by the more subglobose spores and the much more voluminous cells in the pileipellis, additional field differences (and therefore possibly of lesser importance) are the initial brown pigmentation of the cap and stipe fading only later to yellow and the more orange instead of pink color of the hymenophore. Both possess the often laterally compressed, hollow stipe and matted-fibrillose cap surface and are of the same size.

The recently described *Cantharellus conspicuus* Eyssart., Buyck & Verbeken from Zimbabwe (Eyssartier *et al.*, 2002) was also reported as having a general "*Craterellus*" aspect and resembles our species in most of its microscopic features. However, *C. conspicuus* differs markedly in the darker and more purple-lilac colors of its cap and stipe, its squamulose-fibrillose cap and stipe surface, and whitish to pale orange hymenophore.

Other yellow *Cantharellus* species of a similar small size, e.g. *C. microcibarius* Heinem. or *C. isabellinus* var. *parvisporus*, differ by the rarity or absence of clamp connections and their not hollow stipe.

Our species morphologically fits best in subgenus *Parvocantharellus* as defined by Eyssartier & Buyck (2001), being characterized by small basidiomata, abundant clamp connections and absence of thickened cell walls in the hyphal endings of the cap.

**Acknowledgments.** We are grateful to Sida-SAREC through International Science Programme Uppsala University and Molecular Biology project of the University of Dar es Salaam for financial support.

## REFERENCES

- BUYCK B. & NZIGIDAHERA B., 1995 – Ethnomycological notes from western Burundi. *Belgian Journal of Botany* 128: 131–138.
- BUYCK B., EYSSARTIER G., & KIVAISI A., 2000 – Addition to the inventory of the genus *Cantharellus* (Basidiomycota, Cantharellaceae) in Tanzania. *Nova Hedwigia*, 71 (3-4), p. 491-502
- CAMPBELL B., FROST P., & BYRON N., 1996 – Miombo woodlands and their use: overview and key issues. in CAMPBELL, B., (Ed.). *The Miombo in Transition: Woodlands and Welfare in Africa*. CFIOR, Bogor, p. 1-10.
- DANELL E., 1994 – *Cantharellus cibarius: Mycorrhizal formation and ecology*. Universitatis Upsaliensis. Comprehensive Summaries of Uppsala Dissertations from the Faculty of Science and Technology 35, PhD thesis, University of Uppsala, Sweden.

- EYSSARTIER G. & BUYCK B., 2001 — Novitates. Note Nomenclaturale et systematique sur le genre *Cantharellus*. *Documents mycologiques* 121: 55-56.
- EYSSARTIER G., BUYCK B. & VERBEKEN A., 2002 — *Cantharellus conspicuus* sp. nov. *Cryptogamie, Mycologie* 23 (2): 95-102.
- FRIES F., 1821-1832 — *Systema Mycologicum sistens fungorum ordines, genera et species, hucusque cognitatas, quas ad normam methodi naturalis determinavit, disposuit atque descripsit Elias Fries*. Lund (Berlin), Greifswald, 3 vol., 316 p.
- HÄRKÖNEN M., SAARIMAKI T. & MWASUMBI L., 1995 — Edible mushrooms of Tanzania. *Karstenia* 35: 1-92 (supplement).
- HÄRKÖNEN M., NIEMELÄ T. & MWASUMBI L., 2003 — *Tanzanian Mushrooms Edible, harmful and other fungi*. Botanical Museum Finnish Museum of Natural History, 200 pp.
- HEINEMANN P., 1966 — Cantharellineae du Katanga. *Bulletin du Jardin Botanique de l'État*, 36: 365-352
- HÖGBERG P., 1982 — Mycorrhizal associations in some woodlands and forest trees and shrubs in Tanzania. *New Phytologist* 92: 407-415.
- HÖGBERG P., 1992 — Root symbiosis of trees in African dry tropical forests. *Journal of Vegetation Science* 3: 393-400
- HÖGBERG P. & PIEARCE G. D., 1986 — Mycorrhizas in Zambian trees in relation to host taxonomy, vegetation type and successional patterns. *Journal of Ecology* 74: 775-785.
- MORRIS B., 1984 — Macrofungi of Malawi. Some ethnomycological notes. *Bulletin of the British Mycological Society* 18: 45-57.
- MUNYANZIZA E., 1994 — *Miombo trees and mycorrhizae: ecological strategies, a basis for afforestation*. Ph.D. thesis, Wageningen Agricultural University, The Netherlands. 193 p.
- VAN DER MEULEN F., WERGER M.J.A., 1984 — Crown characteristics, leaf size and light through fall of some savanna trees in southern Africa. *South African Journal of Botany* 3 (4): 208-218.

