

## Lichens and lichenicolous fungi new for Tenerife (Canary Islands)

Damien ERTZ<sup>a\*</sup> & Paul DIEDERICH<sup>b</sup>

<sup>a</sup>National Botanical Garden of Belgium, Department of Cryptogamy (BT),  
Domaine de Bouchout, B-1860 Meise, Belgium.  
E-mail: damien.ertz@br.fgov.be

<sup>b</sup>Musée national d'histoire naturelle, 25 rue Munster,  
L-2160 Luxembourg, Luxembourg.  
E-mail: paul.diederich@education.lu

**Résumé** – Une étude de lichens et de champignons lichénicoles de Ténérife, basée sur du matériel récent, est présentée. Deux espèces de lichens sont nouvelles pour l'Afrique : *Anisomeridium ranunculosporum* et *Arthothelium dictyosporum*, ce dernier ayant été considéré jusqu'à présent comme une espèce endémique d'Ecosse. Deux espèces de lichens sont nouvelles pour la Macaronésie : *Brigantiaea leucoxantha* et *B. tricolor*, tandis que la présence de *Polymeridium proponens* y est confirmée. Sept espèces de champignons lichénicoles sont nouvelles pour les îles Canaries : *Lichenosticta alcicorniaria*, *Monodictys epilepraria*, *Paranectria oropensis*, *Phacopsis fusca*, *Phaeosporobolus alpinus*, *Stigmatidium neofusceliae* et *Syzygospora physciacearum*. Quinze espèces sont nouvelles pour Ténérife : *Belonia lumbrispora*, *Biatoropsis usnearum*, *Catillaria nigroclavata*, *Homostegia piggotii*, *Lichenocodium erodens*, *Phacopsis cephalodioides*, *Pronectria pertusariicola*, *Pyrenula hibernica*, *Roselliniella nephromatis*, *Sphaerophorus globosus*, *Syzygospora bachmannii*, *Tremella cetrariicola*, *T. coppinsii*, *T. hypogymniae* et *Xanthoriicola physciae*.

**Biodiversité / biogéographie / hôte / lichen / Macaronésie / Espagne / taxonomie**

**Abstract** – A study of lichens and lichenicolous fungi from Tenerife, based on recent collections, is presented. Two lichen species are new to Africa: *Anisomeridium ranunculosporum* and *Arthothelium dictyosporum*, the latter having been considered until now as an endemic species from Scotland. Two species are new for Macaronesia: *Brigantiaea leucoxantha* and *B. tricolor*, whilst the presence of *Polymeridium proponens* is confirmed. Seven species of lichenicolous fungi are new for the Canary Islands: *Lichenosticta alcicorniaria*, *Monodictys epilepraria*, *Paranectria oropensis*, *Phacopsis fusca*, *Phaeosporobolus alpinus*, *Stigmatidium neofusceliae* and *Syzygospora physciacearum*. Fifteen species are new for Tenerife: *Belonia lumbrispora*, *Biatoropsis usnearum*, *Catillaria nigroclavata*, *Homostegia piggotii*, *Lichenocodium erodens*, *Phacopsis cephalodioides*, *Pronectria pertusariicola*, *Pyrenula hibernica*, *Roselliniella nephromatis*, *Sphaerophorus globosus*, *Syzygospora bachmannii*, *Tremella cetrariicola*, *T. coppinsii*, *T. hypogymniae* and *Xanthoriicola physciae*.

**Biodiversity / biogeography / host / lichen / Macaronesia / Spain / taxonomy**

---

\* Corresponding author.

## INTRODUCTION

Tenerife is the largest and highest island of the Canary Islands and like the rest of the archipelago, is of volcanic origin. The island lies in the mid-Atlantic, between 16.1° and 16.6° W longitudinal and between 28.0° and 28.4° N latitudinal, and has a surface area of around 2000 km<sup>2</sup>. It can be divided in a very dry southern part and a much more humid northern part. The vegetation shows a remarkable altitudinal zonation ranging from the coastal region to the summit of the island at 3718 m (Kunkel, 1993). In the northern part, two large areas (Las Montanas de Anaga and Monte del Agua) are still covered with well preserved laurel forest, which is a subtropical forest type considered as a relic from the Tertiary period. The laurel forest hosts a very rich and original lichen flora such as Lobarion and epiphyllous communities. Many publications were devoted to the flora of lichens and lichenicolous fungi of the Canary Islands archipelago. Most of these data are included in the checklists by Hernández-Padrón (2004) and Hafellner (1995a, 1999, 2002, 2005a). Very recently, contributions to the lichen flora of the archipelago were published on the genera *Usnea* (Clerc 2006) and *Porina* (Sérusiaux *et al.* 2007) and on cyanobacterial lichens (Schultz & van den Boom 2007). New and interesting records were provided for La Palma (van den Boom, 2007). Some new species have also been described from Tenerife (Crespo *et al.*, 2006) and Fuerteventura (van den Boom & Etayo, 2006).

The aim of this paper is to publish additions to the checklist of lichens and lichenicolous fungi of Tenerife, based on the material collected during a field trip of one week made by the two authors. Specimens for this study were collected mainly in coastal areas, laurisilva forests, forests of *Pinus canariensis* and in areas dominated by volcanic rocks with sparse vegetation at different altitudes (see list of localities).

## MATERIAL AND METHODS

Material of lichens and lichenicolous fungi was collected from 18 to 25 July 2007. Examination of thallus and apothecial anatomy was made using standard microscopic techniques. Microscopical examination such as ascospores measurements were made in water and in 5% KOH at magnifications of ×400 or ×1000. The specimens have been deposited in the herbaria of the National Botanical Garden of Belgium (BR) and the private herbarium of Paul Diederich (Di).

### List of Localities

#### Loc. 1.

Tenerife, Las Montanas de Anaga, Monte de las Mercedes, Mirador de Jardina, 28°31' N, 16°17' W, 880 m, trees and rocks along road, 19 and 24 July 2007.

#### Loc. 2.

Tenerife, Las Montanas de Anaga, 3.5 km on road S of Chamorga, 28°33'33" N, 16°09'50" W, 700 m, laurisilva, 19 July 2007.

#### Loc. 3.

Tenerife, S of Puerto de la Cruz, Valle de la Orotava, along road to Teide, between Aguamansa and Bermeja, 28°21'24" N, 16°30'33" W, 1180 m, mixed forest with *Erica arborea* and *Pinus canariensis*, 20 July 2007.

## Loc. 4.

Tenerife, S of Bosque de la Esperanza, along road to Güimar, barranco close to Los Loros, 28°22'18" N, 16°25'39" W, 1125 m, rocks in dry valley, 20 July 2007.

## Loc. 5.

Tenerife, Bosque de la Esperanza, Las Lagunetas, Cumbres de la Victoria, 28°24'34" N, 16°25'16" W, 1675 m, forest with *Pinus canariensis*, 20 July 2007.

## Loc. 6a.

Tenerife, S of Los Silos, c. 1 km W of Erjos, Monte del Agua, 28°19' N, 16°48' W, 1140 m, laurisilva, 22 July 2007.

## Loc. 6b.

Id., forest remnants with *Pinus canariensis*.

## Loc. 6c.

Id., Erjos, on *Casuarina* trees along road (altitude c. 1100 m).

## Loc. 7.

Tenerife, Las Montanas de Anaga, Monte de las Mercedes, Mirador Pico del Inglés, laurisilva and rocks along road, 28°32'00" N, 16°15'51" W, 975 m, 24 July 2007.

## Loc. 8.

Tenerife, Las Montanas de Anaga, Monte de las Mercedes, laurisilva along road and ridge W of Cruz del Carmen, 28°32' N, 16°17' W, 910 m, 24 July 2007.

## Loc. 9.

Tenerife, Las Montanas de Anaga, Monte de las Mercedes, road between Las Mercedes and Mirador de Jardina, 28°31' N, 16°17' W, 860 m, *Cupressus* and other trees along road, 24 July 2007.

## RESULTS

### *Anisomeridium ranunculosporum* (Coppins & P. James) Coppins

Loc. 9, on bark of trees along road, Diederich 16642 & Ertz (Di).

This lichen was previously often misinterpreted as *Arthopyrenia cinereopruinosa* (Coppins, 1988). New for Macaronesia and for Africa.

### *Arthothelium dictyosporum* (Coppins & P. James) Coppins

Loc. 6a, on a smooth trunk in laurisilva, Ertz 10855 (BR).

This species was considered as a possible endemic of Scotland, where it inhabits old woodlands and grows on smooth bark of *Corylus*, *Ilex* and *Sorbus*, as well as on roughish bark of *Salix* (Coppins 1992). It is an extremely rare member of the hyperoceanic *Graphidion* alliance, with previously just six localities known (<http://www.ukbap.org.uk/UKPlans.aspx?ID=113>). As the species is similar in appearance to *Arthonia ilicina* which grows in the same habitats, it might have been overlooked in the laurisilva of Macaronesia where *A. ilicina* is quite common. It is new for Macaronesia and for Africa.

### *Belonia lumbrispora* Etayo

Loc. 8, epiphytic on an old tree in laurisilva, Diederich 16462 & Ertz 10905 (BR, Di).

*B. lumbrispora* is an endemic species of the Canary Islands, described only in 1996 from La Gomera and La Palma (Etayo, 1996a). It is new for Tenerife.

***Biatoropsis usnearum*** Räsänen

Loc. 3, on *Pinus canariensis*, on *Usnea articulata*, Diederich 16700 & Ertz 10792 (BR, Di). Loc. 6a, on trees in laurisilva, on *Usnea*, Diederich 16583 & Ertz 10860, 10870 (BR, Di). Loc. 6b, on *P. canariensis*, on *U. articulata*, Diederich 16648 & Ertz (Di). Monte Aguirre, on *Usnea*, 1990, Becker (Di 9810).

This is a widespread, probably cosmopolitan lichenicolous heterobasidiomycete confined to the host genus *Usnea* s.l. In the Canary Islands, it was known from La Gomera and La Palma (Etayo, 1996b; Hernández-Pradrón, 2004). It is new for Tenerife.

***Brigantiaea leucoxantha*** (Spreng.) R. Sant. & Hafellner

Loc. 2, epiphytic on an old tree in laurisilva, Diederich 16530 & Ertz (Di).

The genus *Brigantiaea* is herewith reported for the first time from Macaronesia. Two species were collected in the same locality. Following Hafellner (1997), one specimen with apothecia reacting only K+ purplish in section belongs to *B. leucoxantha*, whilst other specimens in which the hymenium and hypothecium react K+ bluish violet in addition to the K+ purplish reaction belongs to *B. tricolor*. *B. leucoxantha* is widely distributed in tropical and subtropical regions of Africa and America, whilst the species is rare in Asia and Australia. It is new for Macaronesia. The closest known locality is in Cameroon. Our material from Tenerife has a sorediate thallus, the taxonomic value of which is little understood and regarded as of minor importance by Hafellner (1997).

***Brigantiaea tricolor*** (Mont.) Trevis.

Loc. 2, epiphytic in laurisilva, Diederich 16531 & Ertz 10725 (BR, Di).

This species is widely distributed in the Paleotropics and is also known from Hawaii (Hafellner, 1997). It is new for Macaronesia. The closest known localities are in tropical West Africa (Ivory Coast and Togo). The thallus in our material is sorediate. Following Hafellner (1997), sorediate specimens were known only from the Hawaiian Islands. See also comments under *B. leucoxantha*.

***Catillaria nigroclavata*** (Nyl.) Schuler

Loc. 8, laurisilva along road and ridge, on branch of *Sarothamnus*, Ertz 10911 & Diederich (BR).

In the Canary Islands, this lichen was known from Gran Canaria and La Palma (Hernández-Pradrón, 2004). New for Tenerife.

***Homostegia piggotii*** (Berk. & Broome) P. Karsten

Loc. 6a, on trees in laurisilva, on *Parmelia saxatilis*, Diederich 16596 & Ertz (Di).

In the Canary Islands, this gall-inducing lichenicolous ascomycete was known from La Gomera and La Palma (Hafellner, 1996; Hernández-Pradrón, 2004). It is new for Tenerife.

***Lichenoconium erodens*** M. S. Christ. & D. Hawksw.

Loc. 3, on *Pinus canariensis*, on *Usnea articulata*, Diederich 16701 & Ertz 10782 (BR, Di).

This common and widespread lichenicolous coelomycete was known only from La Palma in the Canary Islands (Hernández-Pradrón, 2004) and is new for Tenerife.

***Lichenosticta alcicorniaria*** (Lindsay) D. Hawksw.

Loc. 6a, on trees in laurisilva, on *Cladonia*, Diederich 16572 & Ertz (Di).

A common lichenicolous coelomycete, usually growing on the lower surface of the squamules of *Cladonia*. New for the Canary Islands.

***Monodictys epilepraria*** Kukwa & Diederich

Loc. 1, on vertical rocks along road, on *Lepraria santosii* (type locality), Diederich 16498 & Ertz (Di). Loc. 8, in laurisilva, on corticolous *Lepraria lobificans*, Diederich 16494 & Ertz (Di).

This lichenicolous fungus has only recently been described from Europe (Kukwa & Diederich, 2005) and it is reported for the first time from the Canary Islands.

***Paranectria oropensis*** (Ces.) D. Hawksw. & Piroz.

Loc. 7, on rocks along road, on *Lecanora*, Diederich 16631 & Ertz 10898 (BR, Di).

This widely distributed lichenicolous fungus is new for the Canary Islands.

***Phacopsis cephalodioides*** (Nyl.) Triebel & Rambold

Loc. 5, on fallen branches on the ground, on *Hypogymnia*, Ertz 10812 & Diederich (BR). Loc. 6a, on tree in laurisilva, on *Hypogymnia*, Ertz 10845 & Diederich (BR).

This lichenicolous ascomycete has rarely been collected but is widely distributed, being known from America (Diederich, 2003), Europe (Triebel *et al.*, 1995) and Russia (Zhurbenko & Otnyukova, 2001). In the Canary Islands, it was known from La Gomera (Hafellner, 1995c) and is here reported as new for Tenerife.

***Phacopsis fusca*** (Triebel & Rambold) Diederich

Loc. 4, on *Xanthoparmelia* with a brown thallus growing on soil over rock, Ertz 10808 p.p. & Diederich (BR, sub *Stigmatidium neofusceliae*).

This widely distributed lichenicolous fungus is new for the Canary Islands.

***Phaeosporobolus alpinus*** R. Sant., Alstrup & D. Hawksw.

Loc. 5, on branches of *Pinus canariensis* in forest, on *Ochrolechia*, Diederich 16663 & Ertz (Di).

*P. alpinus* is a common, probably cosmopolitan lichenicolous fungus on *Ochrolechia*, *Pertusaria* and *Varicellaria* species. It is new for the Canary Islands.

***Polymeridium proponens*** (Nyl.) R. C. Harris

Loc. 2, epiphytic in laurisilva, Diederich 16527 & Ertz 10744 (BR, Di).

Harris (1991) studied material of this lichen species from America, South Africa and India. It has also recently been reported from Tenerife by Hernández-Padrón (2004), but without any references to published data or herbarium material. In the updates to the checklist of insular Macaronesia (Hafellner, 1999, 2002 and 2005a), the species is not reported. The occurrence of the species in Macaronesia is herewith confirmed.

***Pronectria pertusariicola*** Lowen

NE of La Laguna, Anaga-Mountains, Cruz del Carmen, in laurisilva, on *Pertusaria*, 9 April 2003, Diederich 15587 (Di).

In the Canary Islands, this lichenicolous fungus was known from La Gomera (Etayo 1996b). It is new for Tenerife.

***Pyrenula hibernica*** (Nyl.) Aptroot

Loc. 2, small dead standing trunk in laurisilva, Ertz 10757 & Diederich (BR). NE of La Laguna, Anaga-Mountains, Cruz del Carmen, epiphytic in laurisilva, 9 April 2003, Diederich 15586 (Di).

Hernández-Padrón (2004) reported this lichen species from La Gomera and La Palma (as "*Pyrenula chilensis*"). It is therefore new for Tenerife.

***Roselliniella nephromatis*** (Crouan) Matzer & Hafellner

Loc. 6a, on trunk on *Nephroma laevigatum*, Ertz 10833 & Diederich (BR).

This lichenicolous fungus, confined to species of *Nephroma* and *Pseudocyphellaria*, was known so far from Western Europe (Martínez, 2002, Matzer & Hafellner, 1990), Canada (Goward *et al.*, 1996, Etayo & Breuss, 1998) and Macaronesia (Azores, Madeira, La Gomera) (Etayo, 1996b, Kalb & Hafellner, 1992, Kondratyuk & Galloway, 1995). It is new for Tenerife.

***Sphaerophorus globosus*** (Huds.) Vain.

Loc. 6a, epiphytic on tree in laurisilva, Diederich & Ertz (field observation).

In the Canary Islands, this macrolichen was known from El Hierro, La Gomera and La Palma (Hernández-Padrón, 2004). New for Tenerife.

***Stigmatidium neofusceliae*** Calatayud & Triebel

Loc. 4, on *Xanthoparmelia* with a brown thallus growing on earth over rock, Ertz 10808 p. p. & Diederich (BR).

Calatayud & Triebel (1999) recently described this lichenicolous fungus from Spain. Subsequently it has also been reported from the Czech Republic (Kocourková, 2000). New for the Canary Islands.

***Syzygospora bachmannii*** Diederich & M. S. Christ.

Loc. 3, in forest with *Erica arborea*, on terricolous *Cladonia rangiformis*, Diederich 16692 & Ertz (Di); *ibid.*, on trunk of *E. arborea*, on *Cladonia*, Diederich 16685 & Ertz (Di). Loc. 6a, in laurisilva, on *C. furcata*, Diederich 16606 & Ertz (Di). Loc. 8, on terricolous *C. furcata*, Diederich 16472 & Ertz 10910 (BR, Di).

This widely distributed lichenicolous heterobasidiomycete was known from La Palma (Etayo, 2000) and is herewith newly reported from Tenerife.

***Syzygospora physciacearum*** Diederich

Loc. 6c, on *Casuarina* trees along road, on *Physcia adscendens*, Diederich 16568 & Ertz (Di).

This widespread and probably cosmopolitan lichenicolous heterobasidiomycete is here reported as new for the Canary Islands.

***Tremella cetrariicola*** Diederich & Coppins

Loc. 3, on branches in forest with *Erica arborea*, on *Tuckermannopsis chlorophylla*, Diederich 16688 & Ertz 10799 (BR, Di).

In the Canary Islands, this lichenicolous fungus was known from La Gomera (Diederich, 1996; Etayo, 1996b) and La Palma (Etayo, 2000). New for Tenerife.

***Tremella coppinsii*** Diederich & Marson

Loc. 6a, on trees in laurisilva, on *Platismatia glauca*, Diederich 16576 & Ertz 10840 (BR, Di).

In the Canary Islands, this lichenicolous heterobasidiomycete was known from La Gomera (Diederich, 1996; Etayo, 1996b). It is new for Tenerife.

***Tremella hypogymniae*** Diederich & M. S. Christ.

Loc. 6a, on trees in laurisilva, on *Hypogymnia physodes*, Diederich 16594 & Ertz 10826 (BR, Di).

This parasite of *Hypogymnia physodes* is widely distributed in Europe and North America (Diederich, 1996, 2003). In the Canary Islands, it was known from La Gomera (Diederich, 1996; Etayo, 1996b), and is here newly reported from Tenerife.

***Xanthoriicola physciae*** (Kalchbr.) D. Hawksw.

Loc. 6c, on *Casuarina* trees along road, on *Xanthoria parietina*, Diederich 16570 & Ertz (Di).

In the Canary Islands, this lichenicolous hyphomycete confined to *Xanthoria* was known from La Gomera (Hafellner, 1995b) and Lanzarote (Hafellner, 2005b). It is new for Tenerife.

**Acknowledgments.** We would like to thank the “Consejería de Medio Ambiente del Cabildo Insular de Tenerife – Servicio Técnico forestal de Tenerife” for providing us the collecting permit for the protected areas of the island. We are grateful to Brian Coppins for confirming the identification of *Arthothelium dictyosporum* and to André Aptroot for identifying *Polymeridium propionens*.

## REFERENCES

- CALATAYUD V. & TRIEBEL D., 1999 — *Stigmidium neofusceliae* (Dothideales s.l.), a new lichenicolous fungus from Spain. *Nova Hedwigia* 69: 439-448.
- CLERC P., 2006 — Synopsis of *Usnea* (lichenized Ascomycetes) from the Azores with additional information on species in Macaronesia. *Lichenologist* 38: 191-212.
- COPPINS B. J., 1988 — Notes on the genus *Arthopyrenia* in the British Isles. *Lichenologist* 20: 305-325.
- COPPINS B. J., 1992. — *Arthothelium*. In: Purvis *et al.*, The lichen flora of Great Britain and Ireland. London: Natural History Museum Publications: 92-95.
- CRESPO A., ARGÜELLO A., LÜMBSCH H. T., LLIMONA X. & TØNSBERG T., 2006 — A new species of *Lepraria* (Lecanorales: Stereocaulaceae) from the Canary Islands and the typification of *Lepraria isidiata*. *Lichenologist* 38(3): 213-221.
- DIEDERICH P., 1996 — The lichenicolous heterobasidiomycetes. *Bibliotheca Lichenologica* 61: 1-198.
- DIEDERICH P., 2003 — New species and new records of American lichenicolous fungi. *Herzogia* 16: 41-90.
- ETAYO J., 1996a — Contribution to the lichen flora of the Canary Islands. II. Epiphytic lichens from La Palma. *Österreichische Zeitschrift für Pilzkunde* 5: 149-159.
- ETAYO J., 1996b — Aportación a la flora líquénica de las Islas Canarias. I. Hongos liquenícolas de Gomera. *Bulletin de la Société Linnéenne de Provence* 47: 93-110.

- ETAYO J. & BREUSS O., 1998 — New species and interesting records of lichenicolous fungi. *Osterreichische Zeitschrift für Pilzkunde* 7: 203-213.
- ETAYO J., 2000 — Aportación a la flora líquénica de las Islas Canarias. VI. Hongos líquenícolas de La Palma. *Bulletin de la Société linnéenne de Provence* 51: 153-162.
- GOWARD T., DIEDERICH P. & ROSENRETER R., 1996 — Notes on the lichens and allied fungi of British Columbia. III. *The Bryologist* 99: 439-449.
- HAFELLNER J., 1995a — A new checklist of lichens and lichenicolous fungi of insular Laurimacaronesia including a lichenological bibliography for the area. *Fritschiana* 5: 1-135.
- HAFELLNER J., 1995b — Bemerkenswerte Funde von Flechten und lichenicolous Pilzen auf makaronesischen Inseln. I. Erstnachweise einiger Gattungen. In: Daniëls F.J.A., Schulz M., Peine J. (eds.), *Flechten Follmann. Contributions to lichenology in honour of Gerhard Follmann*. Geobotanical and Phytotaxonomical Study Group, Botanical Institute, University of Cologne, Germany, pp. 427-439.
- HAFELLNER J., 1995c — Bemerkenswerte Funde von Flechten und lichenicolous Pilzen auf makaronesischen Inseln III. Einige bisher auf den Kanarischen Inseln übersehene lecanorale Arten. *Linzer Biologische Beiträge* 27: 489-505.
- HAFELLNER J., 1996 — Bemerkenswerte Funde von Flechten und lichenicolous Pilzen auf makaronesischen Inseln IV. Einige bisher übersehene lichenicole Arten der Kanarischen Inseln. *Cryptogamie, Bryologie Lichénologie* 17: 1-14.
- HAFELLNER J., 1997 — A world monograph of *Brigantiaea* (lichenized Ascomycotina, Lecanorales). *Acta Universitatis Upsaliensis, Symbolae Botanicae Upsalienses* 32 (1): 35-74.
- HAFELLNER J., 1999 — Additions and corrections to the checklist and bibliography of lichens and lichenicolous fungi of insular Laurimacaronesia. I. *Fritschiana* 17: 1-26.
- HAFELLNER J., 2002 — Additions and corrections to the checklist and bibliography of lichens and lichenicolous fungi of insular Laurimacaronesia. II. *Fritschiana* 36: 1-10.
- HAFELLNER J., 2005a — Additions and corrections to the checklist and bibliography of lichens and lichenicolous fungi of insular Laurimacaronesia. III. *Fritschiana* 50: 1-13.
- HAFELLNER J., 2005b — Bemerkenswerte Funde von Flechten und lichenicolous Pilzen auf makaronesischen Inseln VII. Über einige bisher auf Lanzarote und Fuerteventura übersehene Arten. *Fritschiana* 50: 15-24.
- HARRIS R.C., 1991 — A revision of *Polymeridium* (Muell. Arg.) R.C. Harris (Trypetheliaceae). *Boletim do Museu Paraense Emílio Goeldi, série Botânica* 7: 619-644.
- HERNÁNDEZ-PADRÓN C. E. (coll. NÚÑEZ S. R., MARTÍN D. S. & VARGAS I. P.), 2004 — Lichens, lichenicolous fungi. In: Izquierdo I., Martín J.L., Zurita N. & Arechavaleta M. (eds), *Lista de especies silvestres de Canarias (hongos, plantas y animales terrestres)*. Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias, pp. 58-82.
- KALB K. & HAFELLNER J., 1992 — Bemerkenswerte Flechten und lichenicole Pilze von der Insel Madeira. *Herzogia* 9: 45-102.
- KOCOURKOVÁ J., 2000 — Lichenicolous fungi of the Czech Republic (the first commented checklist). *Sborník Národního Muzea v Praze, Rada B* 55(3-4): 59-169.
- KONDRATYUK S. & GALLOWAY D. J., 1995 — Lichenicolous fungi and chemical patterns in *Pseudocyphellaria*. *Bibliotheca Lichenologica* 57: 327-345.
- KUKWA M. & DIEDERICH P., 2005 — *Monodictys epilepraria*, a new species of lichenicolous hyphomycetes on *Lepraria*. *Lichenologist* 37: 217-220.
- KUNKEL G., 1993 — Die Kanarischen Inseln und ihre Pflanzenwelt. 3. Auflage. Stuttgart: G. Fischer, 239 pp.
- MARTÍNEZ I., 2002 — Lichenicolous fungi from the Iberian Peninsula and the Macaronesian area. *Nova Hedwigia* 74: 51-67.
- MATZER M. & HAFELLNER J., 1990 — Eine Revision der lichenicolous Arten der Sammelgattung *Rosellinia* (Ascomycetes). *Bibliotheca Lichenologica* 37: 1-138.
- SCHULTZ M. & VAN DEN BOOM P., 2007 — Notes on cyanobacterial lichens (mostly Lichinales, Ascomycota) of the Canary Islands. *Nova Hedwigia* 84: 113-133.
- SÉRUSIAUX E., BERGER F., BRAND M. & VAN DEN BOOM P., 2007 — The lichen genus *Porina* in Macaronesia, with descriptions of two new species. *Lichenologist* 39: 15-33.
- TRIEBEL D., RAMBOLD G. & ELIX J. A., 1995 — A conspectus of the genus *Phacopsis* (Lecanorales). *The Bryologist* 98: 71-83.
- VAN DEN BOOM P. & ETAYO J., 2006 — New records of lichens and lichenicolous fungi from Fuerteventura (Canary Islands), with descriptions of some new species. *Cryptogamie, Mycologie* 27: 341-374.
- VAN DEN BOOM P., 2007 — New and interesting lichenized and lichenicolous fungi from the Canary Island La Palma. *Annalen des Naturhistorischen Museums in Wien* 108B: 153-166.
- ZHURBENKO M. P. & OTNYUKOVA T. N., 2001 — Lichenicolous fungi from the Sayan-Tuvan Mountains, southern Siberia, Russia. *Folia Cryptogamica Estonica* 38: 79-84.