Presence of the foraminifer *Chapmanina gassinensis*
Silvestri, 1931, in the Eocene (Lutetian)
of the Grignon “falunière” (Yvelines, Paris Basin).
The genus *Chapmanina*, its species
and world distribution

Armelle POIGNANT

DIRECTEUR DE LA PUBLICATION : Bruno David,
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Didier Merle

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS*: Emmanuel Côté (geodiv@mnhn.fr); Anne Mabilie

MISE EN PAGE / *PAGE LAYOUT*: Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD*:

Christine Argot (MNHN, Paris)
Beatrix Azanza (Museo Nacional de Ciencias Naturales, Madrid)
Raymond L. Bernor (Howard University, Washington DC)
Alain Blieck (USTL, Villeneuve d'Ascq)
Henning Blom (Uppsala University)
Jean Broutin (UPMC, Paris)
Gaël Clément (MNHN, Paris)
Ted Daeschler (Academy of Natural Sciences, Philadelphie)
Bruno David (MNHN, Paris)
Gregory D. Edgecombe (The Natural History Museum, Londres)
Ursula Göhlich (Natural History Museum Vienna)
Jin Meng (American Museum of Natural History, New York)
Brigitte Meyer-Berthaud (CIRAD, Montpellier)
Zhu Min (Chinese Academy of Sciences, Pékin)
Isabelle Rouget (UPMC, Paris)
Sevket Sen (MNHN, Paris)
Stanislav Štámbek (Museum of Eastern Bohemia, Hradec Králové)
Paul Taylor (The Natural History Museum, Londres)

COUVERTURE / *COVER*:

Réalisée à partir de Figure 1 de l'article/*created from Figure 1 of this article*.

Geodiversitas est indexé dans / *Geodiversitas is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Physical, Chemical, and Earth Sciences®
- Scopus®

Geodiversitas est distribué en version électronique par / *Geodiversitas is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Geodiversitas* sont référencés par /
Articles and nomenclatural novelties published in Geodiversitas are referenced by:

- ZooBank® (<http://zoobank.org>)

Geodiversitas est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Geodiversitas is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:
Adansonia, Zoosystema, Anthropolozologica, European Journal of Taxonomy, Naturae.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle
CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)
Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40
diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2018
ISSN (imprimé / *print*): 1280-9659/ ISSN (électronique / *electronic*): 1638-9395

Presence of the foraminifer *Chapmanina gassinensis* Silvestri, 1931, in the Eocene (Lutetian) of the Grignon “falunière” (Yvelines, Paris Basin). The genus *Chapmanina*, its species and world distribution

Armelle POIGNANT

5 rue Léon Bloy, F-92260 Fontenay-aux-Roses (France)
poignant.armelle@orange.fr

Submitted on 6 July 2017 | accepted on 3 April 2018 | published on 30 August 2018

[urn:lsid:zoobank.org:pub:1336C880-D09D-4C20-8BB1-52B58B378192](https://zoobank.org/pub:1336C880-D09D-4C20-8BB1-52B58B378192)

Poignant A. 2018. — Presence of the foraminifer *Chapmanina gassinensis* Silvestri, 1931, in the Eocene (Lutetian) of the Grignon “falunière” (Yvelines, Paris Basin). The genus *Chapmanina*, its species and world distribution. *Geodiversitas* 40 (17): 461–470. <https://doi.org/10.5252/geodiversitas2018v40a17>. <http://geodiversitas.com/40/17>

KEY WORDS

Foraminifer,
Eocene,
Oligocene,
Paris Basin,
palaeobiogeographical
distribution.

ABSTRACT

Chapmanina gassinensis (foraminifer) has been observed as it seems for the first time in the Lutetian of the “falunière” of Grignon (Yvelines, southwestern Paris Basin). It is widespread in the Eocene of western Europe, also present in the Tethysian domain *l.s.* and even across the Atlantic. In the Oligocene, it becomes very rare and has been only found in some places of Mediterranean Europe and disappears in the Miocene. The Paris Basin seems to be its northernmost occurrence.

RÉSUMÉ

Présence du foraminifère Chapmanina gassinensis Silvestri, 1931, dans l'Éocène (Lutétien) de la falunière de Grignon (Yvelines, Bassin Parisien). Le genre Chapmanina, ses espèces et sa répartition mondiale. Chapmanina gassinensis (foraminifère) a été observée pour la première fois, semble-t-il, dans le Lutétien de la falunière de Grignon (Yvelines, sud-ouest du Bassin de Paris). Elle est largement répandue dans l'Éocène de l'Europe occidentale, présente aussi dans le domaine téthysien *l.s.* et jusque dans le continent américain. À l'Oligocène, elle ne s'observe que dans quelques points de l'Europe méditerranéenne et disparaît ensuite. Le Bassin de Paris apparaît comme sa localisation la plus septentrionale.

MOTS CLÉS

Foraminifère,
Éocène,
Oligocène,
Bassin de Paris,
distribution
paléobiogéographique.

FIG. 1. — *Chapmanina gassinensis* Silvestri, 1931 (Lutetian, Grignon falunière, Yvelines), in lateral (A) and apical (B) views (MNHN.F.F62410). Scale bars: 200 μ m.

INTRODUCTION

Several samples of the Lutetian marl-pit of Grignon (western Paris Basin) have been examined for a research of miliolids. In one of these samples (probably from unit 4, see Fig. 2), I found two specimens of *Chapmanina gassinensis* Silvestri, 1931 (MNHN.F.F62410; Figs 1, 2). This foraminifer is easy to identify and can't be confused with any other one. It is well known in the Eocene of France (Aquitaine Basin) and also observed in the Eocene of various Tethysian European countries and of the Tethysian domain *l.s.* It is only pointed out in the Oligocene in France and a few European countries. Until now, *Ch. gassinensis* had never been mentioned in the Paris Basin.

Sometimes called *Chapmanina* sp. or cf. *gassinensis*, it is very probably *Chapmanina gassinensis* because of the typical associated microfauna. A new species of *Chapmanina* Silvestri, 1931 has been described in the Paleocene of Australia (Quilty & Packham 2006), it will be treated further.

Before studying the genus and its species, it is useful to give some precisions: one of the aims of this work is to show the wide Tethyan distribution of *Chapmanina gassinensis*, but it is probably not an extensive recense of all the localities where this species has been found; in the same way, all the references concerning the localities of its presence in a particular country have not been reported, when they are too many, such as is the case of the Aquitaine in France or of Turkey.

Moreover, it is important to remark that *Chapmanina gassinensis* has not always been considered as larger foraminifera, and so, it has probably been omitted several times in the literature dealing with large benthic assemblages containing for example: *Nummulites*, *Alveolina*, *Discocyclusina*, *Pellatispira*, *Fabiania cassis* (Oppenheim, 1896) with which it is frequently

associated can be up to 4 mm in diameter, *Chapmanina gassinensis* can't exceed 1 mm. According Smout (1954) opinion: "*Chapmanina* has a very small test compared to most complex species".

THE GENUS *CHAPMANINA* AND ITS SPECIES

According Loeblich & Tappan systematics (1987), the genus *Chapmanina* belongs to the family Chapmaninidae Thalmann, 1938, superfamily Rotaliacea Ehrenberg, 1839 and its type species is *Chapmanina gassinensis*, because pointed out at Gassino, in Torino area (Silvestri 1905a). The genus *Chapmanina* first appears with the name *Chapmania* described by Silvestri & Prever (in Silvestri 1904). In 1931, Silvestri changed this name in *Chapmanina*, two species having been described with the name *Chapmania*:

Chapmania galea Silvestri, 1923, type species of the genus *Preverina* Frizzell, 1949, considered by Loeblich & Tappan (1987) as a synonymous of *Chapmanina*, species of the Italian Tortonian (surroundings of Reggio d'Emilia), represented only by an axial section, but the holotype in unknown (Cita & Scipolo 1961), as well as the topotypes.

Chapmania sertata Silvestri, 1929, from the Lutetian of Ancona, Italy, illustrated by a bad section (Frizzell [1949] did not think it is a *Chapmanina*).

Quilty & Packham (2006) described *Chapmanina conjuncta*, in the Australian Paleocene (see further).

In 1918, Halkyard described *Patellina conica* which is in fact *Chapmanina gassinensis*.

Loeblich & Tappan (1987: pl. 775, figs 1-9), reported the genus *Chapmanina* from the Lutetian until the mid-

FIG. 2. — The “Falunière” of Grignon section, after Guernet *et al.* (2012) for the section profile, lithologic units and sub-units and descriptions, Gély (1996) for the sequential units and Huyghe *et al.* (2012) for the correlation of sequential and lithologic units. Section profile modified from Sanders *et al.* 2015.

FIG. 3. — Distribution of *Chapmanina gassinensis* Silvestri, 1931 in the Eocene of Europe and North Africa. The numbers correspond to the references indicated in the Appendix 1.

FIG. 4. — Distribution of *Chapmanina gassinensis* Silvestri, 1931 in the Oligocene of Europe. The numbers correspond to the references indicated in the Appendix 1.

Fig. 5. — Distribution of *Chapmanina* Silvestri, 1931 in the Eocene of Asia. The numbers correspond to the references indicated in the Appendix 1.

dle Miocene in Italia, France, Spain, Greece, Romania. But, I haven't found any available mentions concerning the Miocene, the species *galea* Silvestri, 1923, being quite dubious (see earlier).

Chapmanina gassinensis was emended by Frizzell in 1949 and well studied by Fuchs (1969). Barbin & Decrouez (1987) work is of greatest interest, because they described in detail the complex structure of the species and gave numerous references concerning its distribution. *Chapmanina gassinensis* has been found in the Eocene and the Oligocene associated with a typical fauna of Nummulites and other larger foraminifera: *Nummulites fabianii*, *N. aturicus*, *N. vascus*, *Discocyclina*, *Asterodiscus*, *Assilina*, *Linderina*, *Halkyardia*, *Fabiania*, *Pellatispira*, etc. in Eocene (Fleury *et al.* 1985; Sztrákos & du Fornel 2003); *Nummulites vascus*, *N. intermedius*, *Neorotalia*, *Amphistegina*, etc. in Oligocene (see Boulanger 1968; Cahuzac & Poignant 2002).

THE SPECIES OF THE GRIGNON “FALUNIÈRE” (YVELINES, SOUTH-WESTERN PARIS BASIN)

The lutetian exposure of Grignon called “falunière” is well known in the Paris Basin because of its abundance in various well-preserved organisms (Guernet *et al.* 2012; Sanders *et al.* 2015; Fig. 2). The two found specimens don't allow any doubt about their systematical attribution as their illustrations show (Fig. 1). They have been observed only in one level without any precise localisation, although many

samples have been studied. Le Calvez (1970) did not report it among the numerous listed species, but as it is very rare, it is not surprising. Along my numerous examinations of Grignon sediment, I also met one *Hantkenina* and *Robertina declivis* (Reuss, 1863) which was only reported from the Oligocene.

It is interesting to link the presence of *Chapmanina gassinensis* in the Paris Basin with the one of *Fabiania cassis*. *Fabiania cassis* is mentioned by Bignot (1968) in the upper Lutetian of Mantes (western Paris Basin), then by Le Calvez (1970) in the Lutetian of Paris Basin (Le Tillet drilling), where she saw only a few specimens (cf. also Curry in Le Calvez 1970). Bignot (1968) remarked that *F. cassis*, peri-mediterranean species (Neumann & Boulanger 1955) went rather far to the North during the Eocene. It is the same for *Chapmanina gassinensis*, rather uncommon species, only observed at Grignon until now.

OTHER CITATIONS OF *CHAPMANINA GASSINENSIS* IN FRANCE

Eocene (Fig. 3)

Aquitaine. The Eocene is well represented and has been largely studied. *Ch. gassinensis* has been reported by many authors in numerous localities in the Lutetian, Bartonian, Priabonian. For example: Halkyard (1918: pl. 8, fig. 7): Eocene, blue marl of the “Côte des Basques”, Biarritz (Pyrénées-Atlantiques), named *Patellina* (see above); Cuvillier (1951): Bartonian, *Chapmanina* sp. in Abatilles drill-

ing (Arcachon, south of Bordeaux) and upper Eocene of Biarritz; Sacal & Debourle (1957: pl. 19, fig. 9): middle to upper Eocene of Pontonx (Landes); Magné & Mal-moustier (1964): northern Aquitaine Paleogene; Veillon (1964): lower and middle Eocene of Arcachon; Boulanger (1968: pl. 32, fig. 64; pl. 33, fig. 66): Bartonian, Biarritz and Saint-Cricq-du-Gave (Landes); Deloffre & Hamaoui (1973: pl. 11, fig. 8): upper Eocene of Labatut (Landes); Poignant (1991): upper Eocene of Lac Mouriscot (near Biarritz); Mathelin & Sztrákös (1993): *Truncatulinoidea rohri* zone and lower zone of Bartonian age; Sztrákös *et al.* (1998): Bartonian of Listrac Marls in northern Aquitaine; Sztrákös (2000: pl. 17, fig. 15): Bartonian-Priabonian of Adour valley; Sztrákös *et al.* (2010): very abundant in the upper Eocene of the “Acacias drilling”, near Arcachon.

South-eastern France. Blondeau *et al.* (1968): *Ch. gassinensis* would be abundant in the Alpes-Maritimes, in a transition zone between Lutetian and Upper Eocene; Sztrákös & du Fornel (2003): Bartonian-Priabonian of Alpes-Maritimes and Provence.

Oligocene (Fig. 4)

Aquitaine. *Ch. gassinensis* is rather rarely mentioned in the Oligocene: Veillon (1964): “Les Acacias” drilling; Boulanger (1968): Lower Oligocene, bottom of “Villa Belza” cliff, in Biarritz; Poignant (1972, erroneously attributed to Priabonian), Cahuzac & Poignant (2002): Orx pond, Saint-André-de-Seignanx (Landes).

Cavelier *et al.* (1981), in a publication on the Eocene-Oligocene limit, thought that *Chapmanina gassinensis* disappeared exactly at the Eocene-Oligocene limit. They said: “At the Eocene/Oligocene boundary there was a general drop of temperature accompanied by a period of great drought which was associated with the disappearance or regression of most of the tropical species, for example: *Discocyclina*, *Asterocyclina*, *Lockhartia*, *Fabiania*, *Chapmanina gassinensis*, etc.” (Cavelier *et al.* 1981: 226, 228). This work invalidates this opinion as far as *Chapmanina gassinensis* is concerned.

CITATIONS OF *CHAPMANINA GASSINENSIS* OR Cf. *GASSINENSIS* OR *CHAPMANINA* SP. IN EOCENE OF EUROPE (Fig. 3)

Chapmanina gassinensis has been pointed out in many European countries: Albania, Croatia, Germany, Greece, Hungary, Italy, Spain, Thrace. Frequently named cf. *gassinensis* or *Chapmanina* sp. (for instance: Di Carlo *et al.* (2010), it is without any doubt *gassinensis*, because of the associated microfauna which is the same as in France. Silvestri cited the genus in Toscana (Silvestri 1904), then in Piemonte region (Silvestri 1905a; Gassino, see earlier), in Umbria (Silvestri 1905b) and in Veneto (Silvestri 1923), Romania (Kovacs & Arnaud-Vanneau 2004), Spain (Molina *et al.* 1986) thought that it would be characteristic of the Upper Eocene in Italy and Spain), Thrace (Less *et al.* 2011).

CITATIONS OF *CHAPMANINA GASSINENSIS* OR Cf. *GASSINENSIS* OR *CHAPMANINA* SP. IN OLIGOCENE OF EUROPE (Fig. 4)

Chapmanina gassinensis is reported by Gauthier (1971), in a drilling, in Lluchmajor, in Palma de Mallorca Island; it is frequent and associated with *Lepidocyclinas* and *Miogypsina*s. But this suggests a reworking, since *Chapmanina* has never been observed in the Miocene and B. Cahuzac (pers. comm.) told me that reworking is frequent in Mallorca.

In Italy, Cita & Scipolo (1961) observed *Chapmanina gassinensis* in the Oligocene of Monte Baldo (North-Western Italy), and these authors suggested a reworking; but later on *Chapmanina gassinensis* was found also in the Oligocene of south Apennine and of Vicenza province (Zanfra 1965; de Zanche 1967) suggesting that the occurrence range of this species spans the Eocene to the Oligocene.

Di Carlo *et al.* (2010) mentioned *Chapmanina* sp. in the Rupelian of Zakynthos Island. In Slovenia, Pavlovic *et al.* (1986) found *Chapmanina* sp. in the Oligocene at Bohring and Nova Stifla.

CITATIONS IN TETHYS (MEDITERRANEAN DOMAIN AND EASTERN TETHYS; Fig. 5)

Chapmanines and above all *Chapmanina gassinensis*, sometimes *Chapmanina* sp. have been frequently cited out of Europa, in the Eocene of the Mediterranean domain or further towards East (e.g. in Morocco, Algeria, Tunisia, Syria, Turkey, Oman, Qatar, Tanzania, Iran). In North-eastern India, North of Bangladesh (Matsumaru & Sarma, 2010): *Chapmanina* spp., and *Nummulites atacicus*, *N. globulus*, *N. burdigalensis* and *Fabiania casis* are found in Bartonian. The section figured (Matsumaru & Sarma 2010: pl. 3, fig. 12) does not allow to assert if it is really a *Chapmanina*, but the associated microfauna is of a Bartonian age. The presence of a tethysian microfauna in the East of India corresponds to the persistence of the Tethys ocean which will close at the end of Burdigalian-Ottangien (Rögl 1999). We can also remark the existence of aquitanian faunas (*Miogypsina*, *Lepidocyclina*) in the Qom Basin in Iran (Rahaghi 1973). The rotation of Africa and Arabia meeting the anatolian platform will separate the Mediterranean sea from the Indian Ocean (Rögl 1999).

In Australia, in Adelaïde region, Murray Lindsay & McGowran, 1986, described an Eocene tethysian fauna, with *Halkyardia*, *Linderina*, but they did not mention any *Chapmanina* which could be associated to these species.

In Australia too (off southern New South Wales), Quilty & Packham (2006), described *Chapmanina conjuncta*, in the Paleocene and they saw a clear tethysian influence (see earlier, Murray Lindsay & McGowran 1986). At that period, Tethys was widely opened towards the Indian Ocean. This new species has been observed in a micritic limestone, found at a depth of 1750 m. The specimens are observed in thin sections: “Material: seven thin sections”. Planktonic foraminifera species are very badly preserved and are not easily identifiable as well as calcareous nannoplankton. No *Nummulites* is identified. According to Quilty & Packham (2006), *Chapmanina conjuncta* will differ from *Chapmanina gassinensis* by a larger marginal zone, but their illustration is not conclusive, speci-

mens being known only in thin section. Moreover, the authors say (Quilty & Packham 2006: 331): “In some respects the species found here seems intermediate between *Cymbalopora* van Hagenow, 1851, and *Chapmanina*”. Consequently, it seems difficult to have a reliable opinion. The authors give a Late Paleocene-Early Eocene age to this fauna.

OTHER CITATIONS

Barbin & Decrouez (1987), according to A. Blondeau (pers. comm.), reported the presence of *Chapmanina gassinensis* in the Eocene of Mexico, this is not very surprising since Salmérón (1986) has reported the occurrence of *Fabiania cassis*, often associated with *C. gassinensis*, in the Upper Eocene of Mexico. In the “Foz dos Amazonas Basin”, Central Brasil, de Mello e Souza *et al.* (2003), observed *Fabiania cassis* in the Eocene, associated with american microfaunas. *Chapmanina gassinensis* could also be present. Besides, Fleury *et al.* (1985: fig. 12), have pointed out a current of *Fabiania* coming from Europa and going to America. At last, we can note that the genus *Chapmanina* is present in the Eocene of New-Caledonia (Paris 1981 in Barbin & Decrouez 1987).

CONCLUSION

Chapmanina gassinensis is above all a tethysian affinity species since widespread in the whole Tethys, as well as *Fabiania cassis* to which it appears very often associated. Its distribution is even worldwide since pointed out in the american continent (Upper Eocene of Mexico, see earlier).

We must notice that the extension of *Chapmanina gassinensis* towards the North does not seem to exceed some latitude. The Paris Basin could be its northernmost distribution during the Eocene. It had not cited by Kaasschieter (1961) in his important work on the Eocene of Belgium. Bavaria, where it has been found (Darga 1990) is at a lower latitude.

Very abundant during the Eocene, it becomes rare at the Oligocene, where Greece seems to be its easternmost localisation. It does not occur in the Miocene.

It would be interesting to study the Eocene specimens of *Chapmanina gassinensis* from India since associated to a Mediterranean microfauna: *Nummulites*, *Fabiania*. The presence of the species on the american continent would ask for a work of geographical and stratigraphical distribution, as also the specimen found in New-Caledonia.

The new Australian species, supposed Paleocene, would also require further studies to confirm its age and its taxonomical attribution.

Acknowledgements

I am deeply indebted at first to Annachiara Bartolini who made optical photographs and carefully revised the manuscript and Alexandre Lethiers who realized the outline of the carts of distribution of the species. Special thanks are due to V. Barbin, H. Bismuth, B. Cahuzac, H. Salles-Astier for their bibliographic help and linguistic improvements.

REFERENCES

- AGIP MINERARIA. 1959. — *Microfacies italiane: dal carbonifero al miocene medio*. San Donato milanese, 35 p., 145 pls.
- AKKIRAZ M. S., AKGÜN F., ÖRCEN S., BRUCH A. A. & MOSBRUGGER V. 2006. — Stratigraphic and Paleoenvironmental Significance of Bartonian-Priabonian (Middle-Late Eocene) Microfossils from the Basceme Formation, Denizli Province, Western Anatolia. *Turkish Journal of Earth Sciences* 15: 155-180.
- ALLISON A. 1953. — La geologia della struttura della zona di Priolo (Siracusa). 57^e reunion della Societa Geologica Italiana. Guida alle escursioni, Palermo. *Bollettino della Societa geologica italiana* 71: 109-111.
- ANANIA M. G., BOTTINO C., MATTEUCCI R. & PIGNATTI J. S. 2000. — A late Bartonian-early Priabonian benthonic foraminiferal assemblage from Tornimparte near l'Aquila (Central Italy). *Geologica Romana* 36: 147-161.
- AUBOUIN J. & NEUMANN M. 1959. — Contribution à l'étude stratigraphique et micropaléontologique de l'Éocène en Grèce. *Revue de Micropaléontologie* 2 (1): 31-49.
- AVSAR N. 1991. — Presence of *Nummulites fabianii* (Prever Group (*Nummulites* ex gr. *fabianii*)) and associated foraminifera in the Elazig region. *Mineral research, ExPlate Bulletin* 112: 71-76.
- BABIC L. & ZUPANIC J. 2012. — Laterally variable development of a basin-wide transgressive unit of the North dalmatian foreland basin (Eocene, Dinarides, Croatia). *Geologica croatica* 65 (1): 1-27. <https://doi.org/10.4154/gc.2012.01>
- BARBIN V. & BIGNOT G. 1986. — New proposal for an Eocene-Oligocene boundary according to microfacies from the Priabonian-type section, in POMEROL CH. & PREMOLI-SILVA I. (eds), Terminal Eocene Events. *Developments in Paleontology and Stratigraphy* 9: 49-52. [https://doi.org/10.1016/S0920-5446\(08\)70093-1](https://doi.org/10.1016/S0920-5446(08)70093-1)
- BARBIN V. & DECROUEZ D. 1987. — Le genre *Chapmanina* (foraminifère): État des connaissances et distribution géographique. *Archives des Sciences* 40 (2): 207-224.
- BENEDETTI A. 2010. — Biostratigraphic remarks on the Caltavuturo Formation (Eocene-Oligocene) cropping out at Portella Colla (Madonie Mounts, Sicily). *Revue de Paléontologie de Genève* 29 (1): 197-216.
- BIGNOT G. 1968. — Nouvelle découverte de *Fabiania cassis* (Oppenheim) dans le Lutétien du Bassin de Paris. Colloque sur l'Éocène. Bureau de Recherches géologiques et minières n°58, Paris: 79-81.
- BLONDEAU A., BODELLE J., CAMPREDON R., LANTEAUME M. & NEUMANN M. 1968. — Répartition stratigraphique des grands foraminifères de l'Éocène dans les Alpes-Maritimes (franco-italiennes et Basses-Alpes). Colloque sur l'Éocène. Bureau de Recherches géologiques et minières n°3: 13-26.
- BONNEFOUS J. & BISMUTH H. 1982. — Les faciès carbonatés de plate-forme de l'Éocène moyen et supérieur dans l'offshore tunisien nord-oriental et en mer pélagienne: implications paléogéographiques et analyse micropaléontologique. *Bulletin du Centre de Recherches et d'Exploration-Production, Elf-Aquitaine* 6 (2): 337-403.
- BOULANGER D. 1968. — Révision du Nummulitique de la Chalosse, du Béarn et du Bas-Adour (Landes et Basses-Pyrénées). Thèse de la Faculté des Sciences de l'Université de Paris, 332 p., 41 pls.
- BOZORGNIA F. (AVEC LA COLLABORATION DE BANAFI S.) 1964. — *Microfacies and Microorganisms of Paleozoic through Tertiary Sediments of some parts of Iran*. Natural Iranian oil Company, Teheran-Iran, 22 p., 152 pls.
- CAHUZAC B. & POIGNANT A. 2002. — Associations de foraminifères benthiques dans quelques gisements de l'Oligo-Miocène sud-aquitain. *Revue de Micropaléontologie* 47 (3): 221-256. [https://doi.org/10.1016/S0035-1598\(02\)90027-9](https://doi.org/10.1016/S0035-1598(02)90027-9)
- CASTELLARIN A. & CITA M. B. 1968. — *Chapmanina gassinensis* dans l'Éocène moyen de la coupe de Gallerie. Étude de quelques coupes priaboniennes dans le Monte Baldo (Province Verona et Trento, Italie) et discussions des limites de l'étag. Colloque sur l'Éocène. Bureau de Recherches géologiques et minières n°69: 119-139.

- CAUS E. 1975. — Bioestigrafía del Eoceno medio y superior del Pirineo catalán (y la zona de tránsito entre esta unidad y la cordillera prelitoral catalana). *Revista Espanola de Micropaleontologia* 7 (2): 297-316.
- CAVELIER C., CHATEAUNEUF J. J., POMEROL C., RABUSSIER D., RENARD M. & VERGNAUD-GRAZZINI C. 1981. — The geological events at the Eocene-Oligocene boundary. *Palaeogeography, Palaeoclimatology, Palaeoecology* 36 (3-4): 223-248. [https://doi.org/10.1016/0031-0182\(81\)90108-5](https://doi.org/10.1016/0031-0182(81)90108-5)
- CITA M. B. & SCIPOLLO C. 1961. — *Chapmanina gassinensis* (Silvestri) dans l'Oligocène du Monte Baldo (Italie). *Revue de Micropaléontologie* 4: 121-134.
- CUVILLIER J. (AVEC LA COLLABORATION DE SACAL V.) 1951. — *Correlations stratigraphiques par microfaciès en Aquitaine occidentale*. Leiden, E.J. Brill, 23 p., 90 pls.
- DARGA R. 1990. — The Eisenrichterstein Hallthurm, Bavaria: An upper eocene carbonate ramp (Northern calcareous Alps). *Facies* 23 (1): 17-36. <https://doi.org/10.1007/BF02536705>
- DELOFFRE R. & HAMAOU M. 1973. — Révision des Chapmaninidae et Cymbaloporidae, *Angotia* et *Fabiania* (Foraminifères). *Bulletin du Centre de Recherches de Pau* 7 (2): 291-335.
- DE ZANCHE V. 1967. — Sul ritrovamento di *Chapmanina gassinensis* (Silvestri) nell'Oligocene dei Lessini Orientali (Vicenza). Società Cooperativa Tipografica (Memoria Accademia Patavina 79): 387-390.
- DI CARLO M., ACCORDI G., CARBONE F. & PIGNATTI J. 2010. — Biostratigraphic analysis of Paleogene lowstand wedge conglomerates of a tectonically active platform margin (Zakynthos Island, Greece). *Journal of Mediterranean Earth Sciences* 2: 31-92.
- DURAND-DELGA M. & MAGNÉ J. 1958. — Données stratigraphiques et micropaléontologiques sur le Nummulitique de l'Est des Cordillères bétiques (Espagne). *Revue de Micropaléontologie* 1 (3): 155-175.
- EAMES F. E., BANNER F. T., BLOW W. H. & CLARKE W. J. 1962. — *Fundamentals of Mid-Tertiary Stratigraphic Correlation*. Cambridge University Press, 190 p., 17 pls.
- FERRER J. 1971. — Presencia de macroforaminífera priabonienses en Eocene de Igualada. *Acta geologica hispanica* 6 (1): 4-7.
- FLEURY J. J. 1970. — Le Sénonien et l'Éocène à microorganismes benthoniques du Koklova (zone du Gavrovo, Akarnanie, Grèce continentale). *Revue de Micropaléontologie* 13 (1): 30-44.
- FLEURY J. J., BIGNOT G., BLONDEAU A. & POIGNANT A. 1985. — Biogéographie de foraminifères benthiques téthysiens du Sénonien à l'Éocène supérieur. *Bulletin de la Société géologique de France* 8 (5): 757-770. <https://doi.org/10.2113/gssgfbull.1.5.757>
- FRIZZELL DON L. 1949. — Rotaliid Foraminifera of the Chapmanininae: their natural distinction and parallelism to the *Dictyoconus* lineage. *Journal of Paleontology* 23 (5): 481-495. <https://www.jstor.org/stable/1299475>
- FUCHS H. 1969. — Contributii la Cunoasterea genului *Chapmanina* Silvestri. *Studia Universitatis Babeş-Bolyai, ser. Geologia, Geographia* 14 (1): 61-71.
- GAUTHIER J. C. 1971. — Présence de *Chapmanina gassinensis* (Silvestri, 1905) dans l'Oligocène de l'Île de Palma de Mallorca (Espagne). *Bulletin du Centre de Recherches de Pau* 5 (2): 203-212.
- GÉLY J.-P. 1996. — Le Lutétien du bassin parisien: de l'analyse séquentielle haute résolution à la reconstitution paléogéographique. *Bulletin d'Information des Géologues du Bassin parisien* 34 (2): 3-27.
- GORMUS M., AVSAR N., DINCER F., UYSAL K., KANBUR S., YESILOTKOSE S. & IC Z. 2007. — Benthic foraminiferal biostratigraphy of the Eocene (Lutetian) sediments of the Dinar (Afyon) region. *Bulletin for Earth Sciences* 28 (1): 1-13.
- GUERNET C., HUYGHE D., LARTEAU F., MERLE D., EMMANUEL L., GÉLY J.-P., MICHEL F. & PILET O. 2012. — Les Ostracodes de la falunière de Grignon (Lutétien du Bassin de Paris): implications stratigraphiques. *Geodiversitas* 34 (4): 909-959. <http://dx.doi.org/10.5252/g2012n4a12>
- HALKYARD E. 1918. — *The Fossil Foraminifera of the Blue Marl of the Côtes des Basques, Biarritz*. Notes from the Manchester Museum (Edited with additions by E. Heron-Allen et A. Earland), vol. 24, 145 p.
- HORTSMANN G. 1967. — *Géologie de la partie méridionale de l'île de Zante (Grèce)*. Thèse de Doctorat de l'Université de Paris, 109 p.
- HUYGHE D., MERLE M., LARTEAU F., CHEYPE E. & EMMANUEL L. 2012. — Middle Lutetian climate in the Paris Basin: implications for a marine hotspot of paleobiodiversity. *Facies* 58 (4): 587-604. <https://doi.org/10.1007/s10347-012-0307-3>
- KAASSCHIETER J. P. H. 1961. — *Foraminifera of the Eocene of Belgium*. *Mémoires de l'Institut royal des Sciences naturelles de Belgique*, vol. 147, 271 p.
- KOVACS S. & ARNAUD-VANNEAU A. 2004. — Upper Eocene paleobathymetry approach based on paleoecological assemblages from the Plesca Valley 2; Outcrops, Transylvania. A Preliminary Report. *Acta Paleontologica Romaniaae* 4: 191-202.
- LE CALVEZ Y. 1970. — *Contribution à l'étude des foraminifères paléogènes du Bassin de Paris*. Éditions du Centre national de la Recherche scientifique, Paris (Cahiers de Paléontologie), 326 p.
- LESS G., ÖZCAN & OKAY A. I. 2011. — Stratigraphy and Larger Foraminifera of the Middle Eocene to Lower Oligocene Shallow-Marine Units in the Northern and Eastern Parts of the Thrace Basin, NW Turkey. *Turkish Journal of Earth Sciences* 20: 793-845. <https://doi.org/10.3906/yer-1010-53>
- LOEBLICH A. R. & TAPPAN H. 1987. — *Foraminiferal Genera and their Classification*. Van Nostrand Reinhold Company, New-York, 2 vol., 970 p., 847 pls. <https://doi.org/10.1007/978-1-4899-5760-3>
- LUCIANI V., NEGRI A. & BASSI D. 2002. — The Bartonian-Priabonian transition in the Mossano section (Colli Berici, north-eastern Italy): a tentative correlation between calcareous plankton and shallow-water benthic zonation. *Geobios* 35, suppl. 1: 140-149. [https://doi.org/10.1016/S0016-6995\(02\)00055-4](https://doi.org/10.1016/S0016-6995(02)00055-4)
- MAGNÉ J. & MALMOUSTIER G. 1964. — Contribution à l'étude des associations microfaunistiques du Paléogène nord-aquitain. Colloque sur le Paléogène, Bordeaux, 1954. Bureau de Recherches géologiques et minières, Orléans: 269-282.
- MATHELIN J.-CH. & SZTRÁKOS K. 1993. — *L'Éocène de Biarritz (Pyrénées atlantiques, SW France)*. *Stratigraphie et paléoenvironnements. Monographie des foraminifères*. Éditions du Centre national de la Recherche scientifique, Paris (Cahiers de Micropaléontologie), 85 p.
- MATSUMARU K. & SARMA A. 2010. — Larger foraminiferal biostratigraphy of the Lower Tertiary of Jaintia Hills, Meghalaya, NE India. *Micropaleontology* 56 (6): 539-565.
- MELLO E SOUZA S. H., FAIRCHILD T. R. & TIBANA P. 2003. — Cenozoic biostratigraphy of larger foraminifera from the Foz do Amazonas Basin, Brasil. *Micropaleontology* 49 (3): 253-266. <https://doi.org/10.2113/49.3.253>
- MOLINA E., MONACO P., NOCCHI M. & PARISI G. 1986. — Biostratigraphic correlations between the central subbetic (Spain) and umbro-marchean (Italy) sequences at the Eocene/Oligocene boundary using foraminifera, in POMEROL CH. & PREMOLI-SILVA I. (eds), Terminal Eocene Events. *Developments in Paleontology and Stratigraphy* 9: 75-82. [https://doi.org/10.1016/S0920-5446\(08\)70097-9](https://doi.org/10.1016/S0920-5446(08)70097-9)
- MONTENAT C. & BLONDEAU A. (AVEC LA COLLABORATION DE BIZON G., PERREAU M., RAJU D. S. N. & ROMAN J.) 1977. — Premier aperçu du Tertiaire d'Oman (Péninsule arabique orientale). *Bulletin de la Société géologique de France* 19 (6): 1285-1295. <https://doi.org/10.2113/gssgfbull.19.6.1285>
- MURRAY LINDSAY J. & MCGOWRAN B. 1986. — Eocene-Oligocene boundary in Adelaide region, south Australia, in POMEROL CH. & PREMOLI-SILVA I. (eds), Terminal Eocene Events. *Developments in Paleontology and Stratigraphy* 9: 165-173. [https://doi.org/10.1016/S0920-5446\(08\)70115-8](https://doi.org/10.1016/S0920-5446(08)70115-8)

- NEUMANN M. & BOULANGER D. 1955. — Le genre *Fabiania*. Répartition stratigraphique et géographique en Aquitaine. *Bulletin de la Société géologique de France* 6 (5): 305-311. <https://doi.org/10.2113/gssgfbull.S6-V.4-6.305>
- ORCEN S. 1986. — Biostratigraphy and Paleontology of the Medik-Ebreme (NW Malatya) Area. *Bulletin of the Mineral Research and Exploration* 105 (105-106): 15-45.
- ÖZCAN E., LESS G., OKAY ARAL I., M. BALDI-BECKE, KOLLANYI K. & YILMAZ O. 2010. — Stratigraphy and larger foraminifera of the Eocene shallow-marine and Olistromal units of the Southern Part of the Thrace Basin, NW Turkey. *Turkish Journal of Earth Sciences* 19: 27-77. <https://doi.org/10.3906/yer-0902-11>
- PAVLOVEC R., DROBNE K. & SIKIC L. 1986. — Upper Eocene and Oligocene in Yugoslavia, in POMEROL CH. & PREMOLI-SILVA I. (eds), Terminal Eocene Events. *Developments in Paleontology and Stratigraphy* 9: 109-111. [https://doi.org/10.1016/S0920-5446\(08\)70102-X](https://doi.org/10.1016/S0920-5446(08)70102-X)
- PIERONI P. G. 1965. — *Lepidocyclina* and *Miogypsina* from Opi, Sangro Valley (Central Apennines). *Geologica Romana* 4: 161-180.
- POIGNANT A. 1972. — *Microfaciès et microfaunes du Priabonien, de l'Oligocène et du Miocène d'Aquitaine méridionale*. Université Pierre et Marie Curie, Paris (Travaux du Laboratoire de Micropaléontologie, vol. 1), 11 p.
- POIGNANT A. 1991. — *Unifissurinella*, new genus of microproblematica from the Upper Eocene of southwestern France. *Micro-paleontology* 37 (1): 95-97. <https://doi.org/10.2307/1485748>
- PRIFTY I. & UTA A. 2012. — Relationship of Sazani and Ionian zones based on biostratigraphical data and tectonic facts. *Studia si comunicari. Stiintele Naturi* 28 (2): 193-198.
- QUILTY P. G. & PACKHAM G. H. 2006. — Late Paleocene faunas with *Chapmanina* and *Reticulophragmium* from offshore southeastern Australia: a Tethyan influence. *Alcheringa: an Australian Journal of Paleontology* 30 (2): 315-343. <https://doi.org/10.1080/03115510608619320>
- RAHAGHI A. 1973. — Étude de quelques grands foraminifères de la formation de Qum (Iran central). *Revue de Micropaléontologie* 16 (1): 23-38.
- RÖGL F. 1999. — Mediterranean and Paratethys facts and hypotheses of an Oligocene to Miocene paleogeography (short overview). *Geologica carpathica* 50 (4): 339-349.
- SACAL V. & DEBOURLE A. 1957. — *Foraminifères d'Aquitaine, 2^{ème} partie, Peneroplidae à Victoriellidae*. Mémoires de la Société géologique de France, vol. 78, 88 p.
- SALMERÓN P. 1986. — The Eocene-Oligocene boundary in Mexico, in POMEROL CH. & PREMOLI-SILVA I. (eds), Terminal Eocene Events. *Developments in Paleontology and Stratigraphy* 9: 189-192. [https://doi.org/10.1016/S0920-5446\(08\)70119-5](https://doi.org/10.1016/S0920-5446(08)70119-5)
- SANDERS M. T., MERLE D. & VILLIER L. 2015. — The molluscs of the “Falunière” of Grignon (Middle Lutetian, Yvelines, France): quantification of lithification bias and its impact on the biodiversity assessment of the Middle Eocene of Western Europe. *Geodiversitas* 37 (3): 345-365. <https://doi.org/10.5252/g2015n3a4>
- SCHIAVINOTTO F. 1979. — *Miogypsina* e *Lepidocyclina* del Miocene di Monte La Serra (L'Aquila-Apennino centrale). *Geologica Romana* 18: 253-293.
- SILVESTRI A. 1904. — Località toscana del genera *Chapmania* Silv. et Prev. *Bollettino del Naturalista Siena* 24: 117-119.
- SILVESTRI A. 1905a. — Sul *Dictyoconus aegyptiensis* (Chapman). *Atti della Pontificia Accademia Romana dei Nuovi Lincei* 58: 129-131.
- SILVESTRI A. 1905b. — Notizie sommarie su tre faunule del Lazio Perugia. *Rivista italiana di Paleontologia e Stratigrafia* 11: 140-145.
- SILVESTRI A. 1923. — *Nuovi rinvenimenti di Chapmanie*. *Rendiconti R. Accademia della Società Geologica Scientifica Veneto-Trentino-Istria*, vol. 3, 14: 7-12.
- SILVESTRI A. 1929. — Osservazioni su fossili nummulitica. *Rivista Italiana Paleontologica of Pavia* 35: 15-35.
- SILVESTRI A. 1931. — Sul genere *Chapmanina* e sulla *Alveolina maiellana* n. sp. *Bollettino della Società Geologica italiana* 50: 63-73.
- SINKOVEC B. & SIMUNIC A. 1996. — *The Bauxite Discovered on Mt. Ravna Gora in Hrvatsko Zagorje (Geological Setting, Composition, Origin)*. Rudarsko geoloscà-nafini zbanik, Zagreb, vol. 8: 67-76.
- SIREL E. 1976. — Description of the new species *Eoannularia conica* n. sp. and new observations on the Upper Lutetian-Lower Priabonian limestone in the Darende-Gürün (west of Malatya region). *Bulletin of the Geological Society of Turkey* 19: 79-82.
- SIREL E. & ACAR S. 1993. — *Malatya*, a new foraminiferal genus from the Lutetian of Malatya region (East Turkey). *Geologica Croatica* 46 (2): 181-188. <https://hrcak.srce.hr/19581>
- SMOUT A. H. 1954. — *Lower Tertiary Foraminifera of the Qatar Peninsula*. British Museum of Natural History, London, 96 p.
- SZTRÁKOS K. 1987. — Les foraminifères bartoniens et priaboniens des couches à « *Tritaxia szabo* » de Hongrie et reconstitution paléogéographique de la Montagne de Hongrie au Bartonien et au Priabonien. *Cahiers de Micropaléontologie* 2: 1-37.
- SZTRÁKOS K. 2000. — Les foraminifères de l'Éocène du Bassin de l'Adour (Aquitaine, France): Biostratigraphie et taxinomie. *Revue de Micropaléontologie* 43 (1-2): 71-172.
- SZTRÁKOS K. & DU FERNEL E. 2003. — Stratigraphie, paléocologie et foraminifères du Paléogène des Alpes Maritimes et des Alpes de Haute-Provence (Sud-Est de la France). *Revue de Micropaléontologie* 46: 229-267. <https://doi.org/10.1016/j.revmic.2003.09.003>
- SZTRÁKOS K., BLONDEAU A. & HOTTINGER L. 2010. — Lithostratigraphie et biostratigraphie des formations paléocènes et éocènes nord-aquitaines (bassins de Contis et Parentis, seuil et plate-forme nord-aquitains). Foraminifères éocènes du bassin d'Aquitaine. *Géologie de la France* 2: 3-52.
- SZTRÁKOS K., GÉLY J. P., BLONDEAU A. 1998. — L'Éocène du Bassin sud-aquitain: lithostratigraphie, biostratigraphie et analyse séquentielle. *Géologie de la France* 4: 57-105.
- THIÉBAULT F. 1973. — Étude géologique du Taygète septentrional (Péloponnèse méridional, Grèce). *Annales de la Société Géologique du Nord* 43 (1): 55-74.
- UNGARO S. 1968. — Étude micropaléontologique et stratigraphique de l'Éocène supérieur (Priabonien) de Mossano (Colli Berici). Colloque sur l'Éocène, Paris. Éditions du Bureau de Recherches géologique et minières, vol. 69: 267-281.
- VEILLON M. 1964. — Les zones à foraminifères du Paléogène nord-aquitain et leur valeur stratigraphique. Colloque sur le Paléogène, Bordeaux, 1962, Éditions du Bureau de Recherches géologique et minières, vol. 28: 227-241.
- ZANFRA S. 1965. — Occurrence of *Chapmanina gassinensis* (Silvestri) in the Oligocene of the southern Apennines. *Bollettino delle Società Paleontologica italiana* 4 (2): 302-303.

Submitted on 6 July 2017;
accepted on 3 April 2018;
published on 30 August 2018.

APPENDIX 1. — Numbers mentioned in Figures 1 to 3 attributed to the references of this article.

Reference	Corresp. no. in Figs 3-5	Reference	Corresp. no. in Figs 3-5
Agip mineraria 1959	1	Luciani <i>et al.</i> 2002	42
Akkiraz <i>et al.</i> 2006	2	Magné & Malmoustier 1964	43
Allison 1953	3	Mathelin & Sztrákos 1993	44
Anania <i>et al.</i> 2000	4	Matsumaru & Sarma 2010	45
Aubouin & Neumann 1959	5	Mello e Souza <i>et al.</i> 2003	46
Avsar 1991	6	Molina <i>et al.</i> 1986	47
Babic & Zupanic 2012	7	Montenat & Blondeau 1977	48
Barbin & Bignot 1986	8	Murray Lindsay & McGowran 1986	49
Barbin & Decrouez 1987	9	Neumann & Boulanger 1955	50
Benedetti 2010	10	Orcen 1986	51
Bignot 1968	11	Özcan <i>et al.</i> 2010	52
Blondeau <i>et al.</i> 1968	12	Pavlovic <i>et al.</i> 1986	53
Bonnefous & Bismuth 1982	13	Pieroni 1965	54
Boulanger 1968	14	Poignant 1972	55
Bozorgnia 1964	15	Poignant 1991	56
Cahuzac & Poignant 2002	16	Prifty & Uta 2012	57
Castellarin & Cita 1968	17	Quilty & Packham 2006	58
Caus 1975	18	Rahaghi 1973	59
Cavelier <i>et al.</i> 1981	19	Rögl 1999	60
Cita & Scipolo 1961	20	Sacal & Debourle 1957	61
Cuvillier 1951	21	Salmerón 1986	62
Darga 1990	22	Schiavinotto 1979	63
Deloffre & Hamaoui 1973	23	Silvestri 1904	64
De Zanche 1967	24	Silvestri 1905a	65
Di Carlo <i>et al.</i> 2010	25	Silvestri 1905b	66
Durand-Delga & Magné 1958	26	Silvestri 1923	67
Eames <i>et al.</i> 1962	27	Silvestri 1929	68
Ferrer 1971	28	Silvestri 1931	69
Fleury 1970	29	Sinkovec & Simunic 1996	70
Fleury <i>et al.</i> 1985	30	Sirel 1976	71
Frizzell 1949	31	Sirel & Acar 1993	72
Fuchs 1969	32	Smout 1954	73
Gauthier 1971	33	Sztrákos 1987	74
Gormus <i>et al.</i> 2007	34	Sztrákos 2000	75
Halkyard 1918	35	Sztrákos <i>et al.</i> 2010	76
Hortsmann 1967	36	Sztrákos & du Fornel 2003	77
Kaasschieter 1961	37	Sztrákos <i>et al.</i> 1998	78
Kovacs & Arnaud-Vanneau 2004	38	Thiébaud 1973	79
Le Calvez 1970	39	Ungaro 1968	80
Less <i>et al.</i> 2011	40	Veillon 1964	81
Loeblich & Tappan 1987	41	Zanfra 1965	82